

Waikouaiti Estuary

Broad Scale Habitat Mapping 2016/17



Prepared
for

Otago
Regional
Council

May
2017



Waikouaiti Estuary

Waikouaiti Estuary

Broad Scale Habitat Mapping 2016/17

Prepared for
Otago Regional Council

by

Leigh Stevens and Barry Robertson

Wriggle Limited, PO Box 1622, Nelson 7040, Ph 021 417 936, 0275 417 935, www.wriggle.co.nz



coastalmanagement

iii

RECOMMENDED CITATION:

Stevens, L.M. and Robertson, B.M. 2017. Waikouaiti Estuary: Broad Scale Habitat Mapping 2016/17. Report prepared by Wriggle Coastal Management for Otago Regional Council. 36p.

Contents

Waikouaiti Estuary - Executive Summary	vii
1. Introduction	1
2. Estuary Risk Indicator Ratings.	5
3. Methods	6
4. Results and Discussion	8
4.0. Broad Scale Mapping Summary	8
4.1. Intertidal Substrate (excluding saltmarsh).	8
4.2. Extent of Soft Mud.	10
4.3. Sediment Oxygenation	12
4.4. Opportunistic Macroalgae	12
4.5. Seagrass.	16
4.6. Saltmarsh	18
4.7. 200m Terrestrial Margin	20
4.8. NZ Estuary Trophic Index.	23
5. Summary and Conclusions	24
6. Monitoring Recommendations	25
7. Acknowledgements	26
8. References	26
Appendix 1. Broad Scale Habitat Classification Definitions.	28
Appendix 2. Notes supporting Risk indicator Ratings (Table 2)	29
Appendix 3. Notes on Sampling, Resolution and Accuracy	30
Appendix 4. Opportunistic Macroalgal Blooming Tool	31
Appendix 5. Waikouaiti Estuary Macroalgal Data	35
Appendix 6. Details of Analytical Methods	36

List of Tables

Table 1. Summary of the major environmental issues affecting most New Zealand estuaries.	2
Table 2. Summary of estuary condition risk indicator ratings used in the present report.	5
Table 3. Summary of dominant broad scale features in Waikouaiti Estuary, 2006 and 2016.	8
Table 4. Summary of dominant intertidal substrate, Waikouaiti Estuary, 2016.. . . .	8
Table 5. Grain size results from representative sediments, Waikouaiti Estuary, 2016.. . . .	10
Table 6. Summary of intertidal opportunistic macroalgal cover, Waikouaiti Estuary, Dec. 2016.	14
Table 7. Summary of seagrass (<i>Z. muelleri</i>) cover, Waikouaiti Estuary, 2016.	16
Table 8. Summary of dominant saltmarsh cover, Waikouaiti Estuary, 2016.	18
Table 9. Summary of 200m terrestrial margin land cover, Waikouaiti Estuary, 2016.	20
Table 10. Primary and supporting indicator values used to calculate an ETI score, Waikouaiti Estuary, 2016.	23

List of Figures

Figure 1. Waikouaiti Estuary, showing main estuary zones and fine scale monitoring sites.	4
Figure 2. Visual rating scale for percentage cover estimates of macroalgae (top) and seagrass (bottom). . . .	6
Figure 3. Waikouaiti Estuary - ground-truthing coverage, field photos and location of grain size samples. . .	7
Figure 4. Overview of terrestrial margin and intertidal areas mapped - Waikouaiti Estuary, Dec. 2016. . . .	9
Figure 5. Map of dominant intertidal substrate types - Waikouaiti Estuary, Dec. 2016.	11
Figure 6. Map of areas with low sediment oxygenation - Waikouaiti Estuary, Dec. 2016.	13
Figure 7. Map of intertidal opportunistic macroalgal biomass (g.m ⁻²) - Waikouaiti Estuary, Dec. 2016.. . .	15
Figure 8. Map of intertidal seagrass percentage cover (<i>Zostera muelleri</i>) - Waikouaiti Estuary, Dec. 2016. . .	17
Figure 9. Map of dominant saltmarsh cover - Waikouaiti Estuary, Dec. 2016..	19
Figure 10. Map of 200m Terrestrial Margin - Dominant Land Cover, Waikouaiti Estuary, Dec. 2016.	21
Figure 11. Summary of Catchment Land Cover (LCDB3 2012), Waikouaiti Estuary.	22
Figure A1. Location of macroalgal patches (>5% cover) used in assessing Waikouaiti Estuary, Dec. 2016. . .	35

All photos by Wriggle except where noted otherwise.

WAIKOUAITI ESTUARY - EXECUTIVE SUMMARY

Waikouaiti Estuary is a relatively modified, moderate-large sized (229ha), mesotidal (1.6m spring tidal range), shallow (mean depth ~1-2m at high water), well-flushed (residence time <1 day), seawater-dominated, tidal lagoon type estuary located at Karitane, Otago. It has a single narrow tidal opening that is occasionally restricted, a central river channel, and two main arms. The catchment is dominated by pasture (74%). It is one of the key estuaries in Otago Regional Council's (ORC's) long-term coastal monitoring programme. This report presents the results of the December 2016 broad scale estuary habitat mapping with broad scale monitoring results, overall estuary condition and issues, and monitoring recommendations summarised below.

BROAD SCALE RESULTS

- Intertidal flats comprised 42% of the estuary, saltmarsh 35%, and subtidal waters 23%.
- Intertidal substrates were dominated by firm muddy sand (57%) and very soft mud (19%), with smaller areas of soft muddy sand (9%), firm sand (7%), gravel (4%), soft mud (3%), mobile sand (1%) and artificial boulder fields (<1%).
- Sediment mud content measured within mud habitat was moderate-high (20-29%).
- Opportunistic macroalgal growth (*Ulva intestinalis* and *Gracilaria chilensis*) was sparse overall (~8% of the available intertidal habitat), had an Ecological Quality Rating of "GOOD", and no gross eutrophic zones were present. However, dense macroalgal growths were present in parts of the main channel and subtidally.
- Seagrass (*Zostera muelleri*) covered <2% of the estuary, located in the central basin that has strong tidal flushing.
- Saltmarsh cover was relatively extensive 80ha (45% of the intertidal area) and was dominated by herbfields (96%).
- The 200m terrestrial margin was 75% pasture or unmaintained grassland with 12% densely vegetated buffer.

ESTUARY CONDITION AND ISSUES

In relation to the key issues addressed by the broad scale monitoring (i.e. muddiness, eutrophication, and habitat modification), the 2016 broad scale mapping results show that the estuary supported a variety of substrate types, small areas of high value seagrass beds, and extensive areas of saltmarsh. However it was expressing symptoms of excessive muddiness, a moderate level of eutrophication (i.e. nuisance macroalgal growths present but not combined with soft muds and poor sediment oxygenation in gross eutrophic zones), and historical habitat modification has also been significant.

The combined results place the estuary in a "MODERATE" state overall in relation to ecological health. Fine sediment issues were most evident in the relatively sheltered Merton and East arms, with eutrophication issues previously apparent in these same locations. There has been historically significant modification and loss of estuary saltmarsh (estimated at ~150ha overall with 31ha in the past 10 years) and the 200m terrestrial buffer is now dominated by grassland. Ongoing pressures are evident in these areas.

RECOMMENDED MONITORING

Waikouaiti Estuary has been identified by ORC as a priority for monitoring because it is a moderate-large estuary with high ecological and human use values that is situated in a developed catchment, and therefore vulnerable to excessive sedimentation and eutrophication. Broad scale habitat mapping, in conjunction with fine scale monitoring (including sedimentation rate monitoring), provides valuable information on current estuary condition and trends over time. The following monitoring recommendations are proposed by Wriggle for consideration by ORC. To characterise any issues of change in habitat (e.g. saltmarsh or seagrass area, soft mud extent), it is recommended that broad scale habitat mapping be undertaken at 10 yearly intervals (next scheduled for 2026) unless obvious changes are observed in the interim.

To fully characterise the observed variation in opportunistic macroalgal growth (both green and red blooms) between years, it is recommended that a strong baseline be established by undertaking an annual macroalgal survey (January/February) over the next three years in conjunction with scheduled fine scale sampling.

1. INTRODUCTION

Developing an understanding of the condition and risks to coastal and estuarine habitats is critical to the management of biological resources. The Otago Regional Council's "Regional Policy Statement and Regional Plan: Water" demonstrates the Council's determination to maintain estuaries in good condition. In the period 2005-2008 Otago Regional Council (ORC) undertook preliminary (one-off) monitoring of the condition of seven Otago estuaries in its region. In 2016, ORC began a more comprehensive long-term estuary monitoring programme designed to particularly address the key NZ estuary issues of eutrophication and sedimentation within their estuaries, as well as identifying any toxicity and habitat change issues. The estuaries currently included in the programme are; Shag Estuary, Waikouaiti Estuary and Catlins Estuary.

Monitoring of the Waikouaiti Estuary began with preliminary broad and fine scale monitoring undertaken in November 2006 and the first year of comprehensive baseline monitoring undertaken in December 2016.

Within NZ, the approach for monitoring estuary condition follows the National Estuary Monitoring Protocol (NEMP) (Robertson et al. 2002) and the NZ Estuary Trophic Index (ETI) (Robertson et al. 2016a and b). It consists of three components as follows:

- 1. Ecological Vulnerability Assessment (EVA)** of estuaries in the region to major issues (see Table 1) and appropriate monitoring design. This component has not yet been undertaken on a regional scale for Otago and hence relative vulnerabilities of their estuaries to the key issues have not been formally identified.
- 2. Broad Scale Habitat Mapping (NEMP approach).** This component (see Table 1) maps the key habitats within the estuary, determines their condition, and assesses changes to these habitats over time. Broad scale intertidal mapping of Waikouaiti Estuary was first undertaken in November 2006 (Stewart 2007) and was repeated in December 2016 (Stevens and Robertson 2017). This latter monitoring is the subject of this report.
- 3. Fine Scale Monitoring (NEMP approach).** Monitoring of physical, chemical and biological indicators (see Table 1). This component, which provides detailed information on the condition of Waikouaiti Estuary, was undertaken in a partial form in November 2006 (Stewart 2007), with the first year of baseline monitoring undertaken on 8 December 2016 (Robertson et al. 2017).

Report Structure: The current report presents an overview of key estuary issues in NZ and recommended monitoring indicators (Section 1). This is followed by risk indicator ratings (Section 2) and the sampling methods (Section 3) used in this broad scale assessment. Summarised results of the December 2016 field sampling are then presented and discussed (Section 4) for the following:

- Broad scale mapping of estuary sediment types.
- Broad scale mapping of macroalgal beds (i.e. *Ulva* (sea lettuce), *Gracilaria*).
- Broad scale mapping of seagrass (*Zostera muelleri*).
- Broad scale mapping of saltmarsh vegetation.
- Broad scale mapping of the 200m terrestrial margin surrounding the estuary.

To help the reader interpret the findings, results are related to relevant risk indicator ratings to facilitate the assessment of overall estuary condition (summarised in Section 5), and to guide monitoring recommendations (Section 6).

WAIKOUAITI ESTUARY

Waikouaiti Estuary is a 229ha shallow, intertidal dominated (SIDE) estuary (Figure 1) that discharges via one permanent open (but restricted) tidal mouth to the Pacific Ocean via a broad embayment at Karitane, Otago. Near the sea, a large barrier spit to the north, and the township of Karitane to the south, border the estuary. Further inland it is bordered by farmland, some of which is intensive dairying. Situated at the mouth of the Waikouaiti River (mean flow $\sim 3.5 \text{ m}^3 \cdot \text{s}^{-1}$), the estuary drains a 421km², agriculturally dominated catchment with high producing exotic pastures that support primarily sheep and beef farming occupying 74% of catchment. Some dairy production (total of 750 milking cows) is occurring in the lower catchment bordering the estuary and is aided by irrigation from the main stem of the river (main dairy farm is 513ha, other on Church Rd is 17.8ha). Because the estuary is fed by a relatively small river, the main channel of the upper-mid estuary is poorly flushed during baseflows. As a consequence, this section can become stratified with a surface layer of lighter, low salinity freshwater flowing over a layer of dense saline water at times.

Ecologically, habitat diversity is moderate to high with extensive shellfish beds, large areas of saltmarsh (35% of estuary), unvegetated tidal flats and some seagrass (<1% of estuary). However, the estuary is excessively muddy (22% soft mud), eutrophic in some areas, and the natural vegetated margin has been lost and developed for urban use and grazing. Also, since approximately the 1940's there has been a loss of extensive areas of saltmarsh through drainage and reclamation.

The Waikouaiti Estuary is a high use estuary that is valued for its cultural, spiritual, scientific and aesthetic appeal, its rich biodiversity, shellfish collection, bathing, whitebaiting, fishing, boating, surfing, and walking. The estuary includes a small wharf and moorings for fishing boats. In the "Otago Regional Plan: Water", the Waikouaiti Estuary is listed as a coastal protection area with Kai Tahu cultural and spiritual values. It is also important for coastal birds such as the eastern bar-tailed godwit and oystercatchers.

Table 1. Summary of the major environmental issues affecting most New Zealand estuaries.

1. Sediment Changes

Because estuaries are a sink for sediments, their natural cycle is to slowly infill with fine muds and clays (Black et al. 2013). Prior to European settlement they were dominated by sandy sediments and had low sedimentation rates (<1 mm/year). In the last 150 years, with catchment clearance, wetland drainage, and land development for agriculture and settlements, New Zealand’s estuaries have begun to infill rapidly with fine sediments. Today, average sedimentation rates in our estuaries are typically 10 times or more higher than before humans arrived (e.g. see Abraham 2005, Gibb and Cox 2009, Robertson and Stevens 2007, 2010, and Swales and Hume 1995). Soil erosion and sedimentation can also contribute to turbid conditions and poor water quality, particularly in shallow, wind-exposed estuaries where re-suspension of fine sediments is common. These changes to water and sediment result in negative impacts to estuarine ecology that are difficult to reverse. They include;

- habitat loss such as the infilling of saltmarsh and tidal flats,
- prevention of sunlight from reaching aquatic vegetation such as seagrass meadows,
- increased toxicity and eutrophication by binding toxic contaminants (e.g. heavy metals and hydrocarbons) and nutrients,
- a shift towards mud-tolerant benthic organisms which often means a loss of sensitive shellfish (e.g. pipi) and other filter feeders; and
- making the water unappealing to swimmers.

Recommended Key Indicators:

Issue	Recommended Indicators	Method
Sediment Changes	Soft Mud Area	GIS Based Broad scale mapping - estimates the area and change in soft mud habitat over time.
	Seagrass Area/biomass	GIS Based Broad scale mapping - estimates the area and change in seagrass habitat over time.
	Saltmarsh Area	GIS Based Broad scale mapping - estimates the area and change in saltmarsh habitat over time.
	Mud Content	Grain size - estimates the % mud content of sediment.
	Water Clarity/Turbidity	Secchi disc water clarity or turbidity.
	Sediment Toxicants	Sediment heavy metal concentrations (see toxicity section).
	Sedimentation Rate	Fine scale measurement of sediment infilling rate (e.g. using sediment plates).
Biodiversity of Bottom Dwelling Animals	Type and number of animals living in the upper 15cm of sediments (infauna in 0.0133m ² replicate cores), and on the sediment surface (epifauna in 0.25m ² replicate quadrats).	

2. Eutrophication

Eutrophication is a process that adversely affects the high value biological components of an estuary, in particular through the increased growth, primary production and biomass of phytoplankton, macroalgae (or both); loss of seagrass, changes in the balance of organisms; and water quality degradation. The consequences of eutrophication are undesirable if they appreciably degrade ecosystem health and/or the sustainable provision of goods and services (Ferriera et al. 2011). Susceptibility of an estuary to eutrophication is controlled by factors related to hydrodynamics, physical conditions and biological processes (National Research Council, 2000) and hence is generally estuary-type specific. However, the general consensus is that, subject to available light, excessive nutrient input causes growth and accumulation of opportunistic fast growing primary producers (i.e. phytoplankton and opportunistic red or green macroalgae and/or epiphytes - Painting et al. 2007). In nutrient-rich estuaries, the relative abundance of each of these primary producer groups is largely dependent on flushing, proximity to the nutrient source, and light availability. Notably, phytoplankton blooms are generally not a major problem in well flushed estuaries (Valiela et al. 1997), and hence are not common in the majority of NZ estuaries. Of greater concern are the mass blooms of green and red macroalgae, mainly of the genera *Cladophora*, *Ulva*, and *Gracilaria* which are now widespread on intertidal flats and shallow subtidal areas of nutrient-enriched New Zealand estuaries. They present a significant nuisance problem, especially when loose mats accumulate on shorelines and decompose, both within the estuary and adjacent coastal areas. Blooms also have major ecological impacts on water and sediment quality (e.g. reduced clarity, physical smothering, lack of oxygen), affecting or displacing the animals that live there (Anderson et al. 2002, Valiela et al. 1997).

Recommended Key Indicators:

Issue	Recommended Indicators	Method
Eutrophication	Macroalgal Cover/Biomass	Broad scale mapping - macroalgal cover/biomass over time.
	Phytoplankton (water column)	Chlorophyll a concentration (water column).
	Sediment Organic and Nutrient Enrichment	Chemical analysis of sediment total nitrogen, total phosphorus, and total organic carbon concentrations.
	Water Column Nutrients	Chemical analysis of various forms of N and P (water column).
	Redox Profile	Redox potential discontinuity profile (RPD) using visual method (i.e. apparent Redox Potential Depth - aRPD) and/or redox probe. Note: Total Sulphur is also currently under trial.
	Biodiversity of Bottom Dwelling Animals	Type and number of animals living in the upper 15cm of sediments (infauna in 0.0133m ² replicate cores), and on the sediment surface (epifauna in 0.25m ² replicate quadrats).

Table 1. Summary of major environmental issues affecting New Zealand estuaries (continued).

3. Disease Risk

Runoff from farmland and human wastewater often carries a variety of disease-causing organisms or pathogens (including viruses, bacteria and protozoans) that, once discharged into the estuarine environment, can survive for some time (e.g. Stewart et al. 2008). Every time humans come into contact with seawater that has been contaminated with human and animal faeces, we expose ourselves to these organisms and risk getting sick. Human diseases linked to such organisms include gastroenteritis, salmonellosis and hepatitis A (Wade et al. 2003). Aside from serious health risks posed to humans through recreational contact and shellfish consumption, pathogen contamination can also cause economic losses due to closed commercial shellfish beds.

Recommended Key Indicators:

Issue	Recommended Indicators	Method
Disease Risk	Shellfish and Bathing Water faecal coliforms, viruses, protozoa etc.	Bathing water and shellfish disease risk monitoring (Council or industry driven).

4. Toxic Contamination

In the last 60 years, NZ has seen a huge range of synthetic chemicals introduced to the coastal environment through urban and agricultural storm-water runoff, groundwater contamination, industrial discharges, oil spills, antifouling agents, leaching from boat hulls, and air pollution. Many of them are toxic even in minute concentrations, and of particular concern are polycyclic aromatic hydrocarbons (PAHs), heavy metals, polychlorinated biphenyls (PCBs), endocrine disrupting compounds, and pesticides. When they enter estuaries these chemicals collect in sediments and bio-accumulate in fish and shellfish, causing health risks to marine life and humans. In addition, natural toxins can be released by macroalgae and phytoplankton, often causing mass closures of shellfish beds, potentially hindering the supply of food resources, as well as introducing economic implications for people depending on various shellfish stocks for their income. For example, in 1993, a nationwide closure of shellfish harvesting was instigated in NZ after 180 cases of human illness following the consumption of various shellfish contaminated by a toxic dinoflagellate, which also led to wide-spread fish and shellfish deaths (de Salas et al. 2005). Decay of organic matter in estuaries (e.g. macroalgal blooms) can also cause the production of sulphides and ammonia at concentrations exceeding ecotoxicity thresholds.

Recommended Key Indicators:

Issue	Recommended Indicators	Method
Toxins	Sediment Contaminants	Chemical analysis of heavy metals (total recoverable cadmium, chromium, copper, nickel, lead and zinc) and any other suspected contaminants in sediment samples.
	Biota Contaminants	Chemical analysis of suspected contaminants in body of at-risk biota (e.g. fish, shellfish).
	Biodiversity of Bottom Dwelling Animals	Type and number of animals living in the upper 15cm of sediments (infauna in 0.0133m ² replicate cores), and on the sediment surface (epifauna in 0.25m ² replicate quadrats).

5. Habitat Loss

Estuaries have many different types of high value habitats including shellfish beds, seagrass meadows, saltmarshes (rushlands, herbfields, reedlands etc.), tidal flats, forested wetlands, beaches, river deltas, and rocky shores. The continued health and biodiversity of estuarine systems depends on the maintenance of high-quality habitat. Loss of such habitat negatively affects fisheries, animal populations, filtering of water pollutants, and the ability of shorelines to resist storm-related erosion. Within New Zealand, habitat degradation or loss is common-place with the major causes being sea level rise, population pressures on margins, dredging, drainage, reclamation, pest and weed invasion, reduced flows (damming and irrigation), over-fishing, polluted runoff, and wastewater discharges (IPCC 2007 and 2013, Kennish 2002).

Recommended Key Indicators:

Issue	Recommended Indicators	Method
Habitat Loss	Saltmarsh Area	Broad scale mapping - estimates the area and change in saltmarsh habitat over time.
	Seagrass Area	Broad scale mapping - estimates the area and change in seagrass habitat over time.
	Vegetated Terrestrial Buffer	Broad scale mapping - estimates the area and change in buffer habitat over time.
	Shellfish Area	Broad scale mapping - estimates the area and change in shellfish habitat over time.
	Unvegetated Habitat Area	Broad scale mapping - estimates the area and change in unvegetated habitat over time, broken down into the different substrate types.
	Sea level	Measure sea level change.
	Others e.g. Freshwater Inflows, Fish Surveys, Floodgates, Wastewater Discharges	Various survey types.

1. INTRODUCTION (CONTINUED)



Figure 1. Waikouaiti Estuary, showing main estuary zones and fine scale monitoring sites.

2. ESTUARY RISK INDICATOR RATINGS

The estuary monitoring approach used by Wriggle has been established to provide a defensible, cost-effective way to help quickly identify the likely presence of the predominant issues affecting NZ estuaries (i.e. eutrophication, sedimentation, disease risk, toxicity and habitat change; Table 1), and to assess changes in the long term condition of estuarine systems. The design is based on the use of primary indicators that have a documented strong relationship with water or sediment quality.

In order to facilitate this assessment process, “risk indicator ratings” have also been proposed that assign a relative level of risk (e.g. very low, low, moderate, high) of specific indicators adversely affecting intertidal estuary condition (see Table 2 below). Each risk indicator rating is designed to be used in combination with relevant information and other risk indicator ratings, and under expert guidance, to assess overall estuarine condition in relation to key issues, and make monitoring and management recommendations. When interpreting risk indicator results we emphasise:

- The importance of taking into account other relevant information and/or indicator results before making management decisions regarding the presence or significance of any estuary issue e.g. community aspirations, cost/benefit considerations.
- That rating and ranking systems can easily mask or oversimplify results. For instance, large changes can occur within the same risk category, but small changes near the edge of one risk category may shift the rating to the next risk level.
- Most issues will have a mix of primary and supporting indicators, primary indicators being given more weight in assessing the significance of results. It is noted that many supporting estuary indicators will be monitored under other programmes and can be used if primary indicators reflect a significant risk exists, or if risk profiles have changed over time.
- Ratings have been established in many cases using statistical measures based on NZ estuary data and presented in the NZ estuary Trophic Index (NZ ETI; Robertson et al. 2016a and 2016b). However, where such data is lacking, or has yet to be processed, ratings have been established using professional judgement, based on our experience from monitoring numerous NZ estuaries. Our hope is that where a high level of risk is identified, the following steps are taken:
 1. Statistical measures be used to refine indicator ratings where information is lacking.
 2. Issues identified as having a high likelihood of causing a significant change in ecological condition (either positive or negative), trigger intensive, targeted investigations to appropriately characterise the extent of the issue.
 3. The outputs stimulate discussion regarding what an acceptable level of risk is, and how it should best be managed.

The indicators and interim risk ratings used for the Waikouaiti Estuary broad scale monitoring programme are summarised in Table 2, with supporting notes explaining the use and justifications for each indicator in Appendix 2. The basis underpinning most of the ratings is the observed correlation between an indicator and the presence of degraded estuary conditions from a range of tidal lagoon and tidal river estuaries throughout NZ. Work to refine and document these relationships is ongoing.

Table 2. Summary of estuary condition risk indicator ratings used in the present report.

RISK INDICATOR RATINGS / ETI BANDS (indicate risk of adverse ecological impacts)				
BROAD AND FINE SCALE INDICATORS	Very Low - Band A	Low - Band B	Moderate - Band C	High - Band D
Soft mud (% of unvegetated intertidal substrate)*	<1%	1-5%	>5-15%	>15%
Sediment Mud Content (%mud)*	<5%	5-10%	>10-25%	>25%
Apparent Redox Potential Discontinuity (aRPD)**	Unreliable	Unreliable	0.5-2cm	<0.5cm
Redox Potential (RP mV) upper 3cm***	>+100mV	+100 to -50mV	-50 to -150mV	<-150mV
Sediment Oxygenation (aRPD <0.5cm or RP@3cm <-150mV)*	<0.5ha or <1%	0.5-5ha or 1-5%	6-20ha or >5-10%	>20ha or >10%
Macroalgal Ecological Quality Rating (OMBT)*	≥0.8 - 1.0	≥0.6 - <0.8	≥0.4 - <0.6	0.0 - <0.4
Seagrass (% change from baseline)	<5% decrease	5%-10% decrease	>10-20% decrease	>20% decrease
Gross Eutrophic Zones (ha or % of intertidal area)	<0.5ha or <1%	0.5-5ha or 1-5%	6-20ha or >5-10%	>20ha or >10%
Saltmarsh Extent (% of intertidal area)	>20%	>10-20%	>5-10%	0-5%
Supporting indicator Extent (% remaining from est. natural state)	>80-100%	>60-80%	>40-60%	<40%
Vegetated 200m Terrestrial Margin	>80-100%	>50-80%	>25-50%	<25%
Percent Change from Monitored Baseline	<5%	5-10%	>10-20%	>20%
NZ ETI score*	0-0.25	0.25-0.50	0.50-0.75	0.75-1.0

*NZ ETI (Robertson et al. 2016b), **Hargrave et al. (2008), ***Robertson (PhD in prep.), Keeley et al. (2012). See NOTES in Appendix 2 for further information.

3. METHODS

Broad-scale mapping is a method for describing habitat types based on the dominant surface features present (e.g. substrate: mud, sand, cobble, rock; or vegetation: macrophyte, macroalgae, rush-land, etc). It follows the NEMP approach originally described for use in NZ estuaries by Robertson et al. (2002) with a combination of detailed ground-truthing of aerial photography, and GIS-based digital mapping from photography to record the primary habitat features present. Appendix 1 lists the definitions used to classify substrate and saltmarsh vegetation. Very simply, the method involves:

- Obtaining aerial photos of the estuary for recording dominant habitat features.
- Carrying out field identification and mapping (i.e. ground-truthing) using laminated aerial photos.
- Digitising ground-truthed features evident on aerial photographs into GIS layers (e.g. ArcMap).

The georeferenced spatial habitat maps provide a robust baseline of key indicators that are used with risk ratings to assess estuary condition in response to common stressors, and assess future change.

Estuary boundaries were set seaward from an imaginary line closing the mouth to the upper extent of saline intrusion (i.e. where ocean derived salts during average annual low flow are <0.5ppt). For the current study, LINZ rectified colour aerial photos (~0.25m/pixel resolution) flown in 2014/15 were sourced from ESRI online, laminated (scale of 1:3,000), and used by experienced scientists who walked the area in Dec. 2016 to ground-truth the spatial extent of dominant vegetation and substrate types. From representative broad scale substrate classes, 6 grain size samples were analysed to validate substrate classifications (Figure 3, Table 5). When present, macroalgae and seagrass patches were mapped to the nearest 5% using a 6 category percent cover rating scale as a guide to describe density (see Figure 2). Notes on sampling, resolution and accuracy are presented in Appendix 3.


Macroalgae was further assessed by identifying patches of comparable growth, and enumerating each patch by measuring:

- % cover of opportunistic macroalgae (the spatial extent and density of algal cover providing an early warning of eutrophication issues).
- macroalgal biomass (providing a direct measure of areas of excessive growth).
- extent of algal entrainment in sediment (highlighting where nuisance condition have a high potential for establishing and persisting).
- gross eutrophic zones (highlighting significant sediment degradation by measuring where there is a combined presence of high algal cover or biomass, low sediment oxygenation, and soft muds).

Where macroalgal cover exceeded 5% of the Available Intertidal Habitat (AIH), a modified Opportunistic Macroalgal Blooming Tool (OMBT) is used to rate macroalgal condition. The OMBT, described in detail in Appendix 2, is a 5 part multimetric index that produces an overall Ecological Quality Rating (EQR) ranging from 0 (major disturbance) to 1 (minimally disturbed) and which is placed within overall quality status threshold bands (i.e. bad, poor, good, moderate, high - Appendix 2). This integrated index provides a comprehensive measure of the combined influence of macroalgal growth and distribution.

Broad scale habitat features were digitised into ArcMap 10.5 shapefiles using a Wacom Cintiq21UX drawing tablet, and combined with field notes and georeferenced photographs, to produce habitat maps showing the dominant cover of: substrate, macroalgae (e.g. *Ulva*, *Gracilaria*), saltmarsh vegetation, and the 200m wide terrestrial margin vegetation/landuse (Figure 4). These broad scale results are summarised in Section 4, with the supporting GIS files (supplied as a separate electronic output) providing a much more detailed data set designed for easy interrogation to address specific monitoring and management questions.

Figure 2. Visual rating scale for percentage cover estimates of macroalgae (top) and seagrass (bottom).



3. METHODS (CONTINUED)



Figure 3. Waikouaiti Estuary - mapped estuary extent showing ground-truthing coverage, field photos and location of grain size samples used to validate substrate classifications.

4. RESULTS AND DISCUSSION

4.0. BROAD SCALE MAPPING SUMMARY

The 2016 broad scale habitat mapping ground-truthed and mapped all intertidal substrate and vegetation including the dominant land cover of the 200m terrestrial margin, with the six dominant estuary features summarised in Table 3 and shown in Figure 4. The estuary comprises an enclosed tidal estuary dominated by intertidal flats (42%) and saltmarsh (35%) that is located predominantly within sheltered tidal arms present on both sides of the main subtidal river channel (23%). Intertidal seagrass (<1%) was sparse, as was dense (>50% cover) opportunistic macroalgae (2%). The 200m wide terrestrial margin was dominated by grassland (75%) with just 12% densely vegetated buffer. The supporting GIS files underlying this written report provide a detailed spatial record of the key features present throughout the estuary. These are intended as the primary supporting tool to help the Council address a wide suite of estuary issues and management needs, and to act as a baseline to assess future change.

While mapping of the estuary was previously undertaken in 2006 (Stewart 2007), the supplied GIS files included only high level summary detail which limited the extent that detailed comparisons could be made. The 2006 and 2016 mapping results are summarised in Table 3.

In the following sections, various factors related to each of these key habitats (e.g. area of soft mud) are used in conjunction with risk ratings to assess key estuary issues of sedimentation, eutrophication, and habitat modification.

Table 3. Summary of dominant broad scale features in Waikouaiti Estuary, 2006 and 2016.

Dominant Estuary Feature	2006			2016		
	ha	% intertidal	% estuary	ha	% intertidal	% estuary
1. Intertidal flats (excluding saltmarsh)	87	44	35	96.6	54.6	42.2
2. Opportunistic macroalgal beds (>50% cover) [on intertidal flats]	0.6	0.3	0.2	5.2	2.9	2.3
3. Seagrass (>20% cover) [on intertidal flats]	0.2	0.1	0.1	1.6	0.9	0.7
4. Saltmarsh	111	56	44	80.3	45.4	35.1
5. Subtidal waters (not recorded in 2006, 2016 value used)	52 ¹	-	21	52.2	-	22.8
Total Estuary	251	100%		229		100
6. 200m wide densely vegetated Terrestrial Margin (e.g. scrub, shrub, forest)						12%

4.1. INTERTIDAL SUBSTRATE (EXCLUDING SALTMARSH)

Results (summarised in Table 4 and Figure 5) show substrates on intertidal flats were dominated by firm muddy sand (57%) and very soft mud (19%), with smaller areas of soft muddy sand (9%), firm sand (7%), gravel (4%), soft mud (3%), mobile sand (1%) and artificial boulder fields (<1%). Strong tidal and freshwater flushing action in the main river channel facilitates the removal of fine material and helps to maintain gravel beds in the upper estuary, and sand dominated substrate in the lower estuary. In the more sheltered arms either side of the main channel, where current flows are much less pronounced, muddy sands dominate in the areas closest to the main channel, while soft and very soft muds are confined largely to the upper tidal reaches.

Table 4. Summary of dominant intertidal substrate, Waikouaiti Estuary, 2016.

Dominant Substrate	Ha	%	Comments
Built feature	0.4	0.4	Artificial boulder fields along reclamation margins
Gravel field	3.4	3.5	Upper estuary river margins
Mobile sand	1.3	1.4	Predominantly near the entrance
Firm sand	6.9	7.1	Small pockets in the main basin and near the entrance
Soft muddy sand	8.7	9.0	Within the sheltered basins of the Merton and East arms
Firm muddy sand	55.0	56.9	Main river channel margins and lower settlement basins
Soft mud	2.7	2.8	Mid to upper settlement basins
Very soft mud	18.2	18.8	Upper settlement basins and sheltered channels
Grand Total	96.6	100	

4. RESULTS AND DISCUSSION (CONTINUED)




Figure 4. Overview of terrestrial margin and intertidal areas mapped - Waikouaiti Estuary, Dec. 2016.

4. RESULTS AND DISCUSSION (CONTINUED)

4.2. EXTENT OF SOFT MUD

Where soil erosion from catchment disturbance exceeds the assimilative capacity of an estuary, adverse estuary impacts are expected from increased muddiness and turbidity, shallowing, increased nutrients, increased organic matter degradation by anoxic processes (e.g. sulphide production), increased contaminant concentrations (where fine muds provide a sink for catchment contaminants like heavy metals), and alterations to saltmarsh, seagrass, fish and invertebrate communities. In particular, multiple studies have shown estuarine macroinvertebrate communities to be adversely affected by mud accumulation, both through direct and indirect mechanisms including: declining sediment oxygenation, smothering, and compromise of feeding habits (e.g. see Mannino and Montagna 1997; Rakocinski et al. 1997; Peeters et al. 2000; Norkko et al. 2002; Ellis et al. 2002; Thrush et al. 2003; Lohrer et al. 2004; Sakamaki and Nishimura 2009; Wehkamp and Fischer 2012; Robertson 2013).

Because of such consequences, three key measures are commonly used to assess soft mud:

- i. **Horizontal extent** (area of soft mud) - broad scale indicator (see rating in Table 2)
- ii. **Vertical buildup** (sedimentation rate) - fine scale assessment using sediment plates (or retrospectively through historical coring). Ratings are currently under development as part of national ANZECC guidelines.
- iii. **Sediment mud content** - fine scale indicator - recommended guideline is no increase from established baseline.

The area (horizontal extent) of intertidal soft mud is the primary sediment indicator used in the current broad scale report, with sediment mud content a supporting indicator. Sediment plates have been established at two fine scale sites to enable future monitoring of vertical buildup (see Robertson et al. 2017 for details).

Figure 5 and Table 4 shows that soft or very soft muds covered 20.9ha (21.7%) of the intertidal area, a risk indicator rating of HIGH, and had a mud content measured in representative areas of 20-29%, a supporting risk indicator rating of MODERATE to HIGH (Table 5). Soft muds were concentrated in the upper tidal reaches of the estuary embayments where mud settlement is thought to predominantly reflect the presence of sheltered deposition zones and, to a lesser extent, salinity driven flocculation. Within the dominant firm sandy mud substrate alongside the main channel, grain size reflected a LOW risk rating (5-10% mud content).

The 2016 soft mud extent (20.9ha) was approximately half that reported in 2006 (39.1ha, Stewart 2007), with much less mud evident in the Merton Arm in 2016 than previously recorded.



Table 5. Grain size results from representative sediments, Waikouaiti Estuary, 2016.

Broad Scale Classification	Site #	% mud	% sand	% gravel	NZTM East	NZTM North
Firm Muddy Sand	6	5.9	77.4	16.7	1416918	4945092
Firm Muddy Sand	5	8.6	91	0.4	1416318	4945228
Soft Mud	1	20.2	69.4	10.4	1416960	4945588
Soft Mud	2	25.2	71	3.9	1416351	4945244
Very Soft Mud	4	27.6	71.7	0.7	1417076	4945821
Very Soft Mud	3	28.6	70.8	0.5	1415687	4945184

See Figure 3 for site locations

4. RESULTS AND DISCUSSION (CONTINUED)

Substrate Class

-  Built feature
-  Rock field
-  Boulder field
-  Cobble field
-  Gravel field
-  Shell bank
-  Mobile sand
-  Firm sand
-  Firm muddy sand
-  Firm mud
-  Soft sand
-  Soft muddy sand
-  Soft mud
-  Very soft mud
-  Water



Figure 5. Map of dominant intertidal substrate types - Waikouaiti Estuary, Dec. 2016.

4. RESULTS AND DISCUSSION (CONTINUED)

4.3. SEDIMENT OXYGENATION

The primary indicators used to assess sediment oxygenation are aRPD depth and RP measured at 3cm. These indicators were measured at representative sites throughout the dominant sand and mud substrate types, and from a range of sites with variable macroalgal cover and bio-mass. From these measurements, broad boundaries have been drawn of estuary zones where sediment oxygen is depleted to the extent that adverse impacts to macrofauna (sediment and surface dwelling animals) are expected (Figure 6). Because macrofauna are used as an indicator of ecological impacts to other taxa, it is expected that these zones will also be exerting adverse impacts on associated higher trophic communities including birds and fish.

These results show that the majority of the estuary sediments are well to moderately well oxygenated and appeared in good (healthy) ecological condition, with the aRPD depth at 2-5cm and the RP above -150mV at 3cm in most sand and gravel dominated sediments. However, there was also 23.5ha (10%) of the total intertidal area identified as having depleted sediment oxygen, a NZ ETI risk rating of "HIGH". These areas were largely confined to soft and very soft muds located in the upper tidal range settlement basins in the Merton and East arms. In 2013 these areas supported dense growths of the red nuisance macroalgae *Gracilaria* that had smothered the estuary bed. While nuisance macroalgae was not present in December 2016, the impact of the previous growth, and its subsequent degradation, is evident in the organically enriched and oxygen depleted sediments that remain in these areas. Sediment oxygenation was not recorded outside of fine scale sites in 2006.

4.4. OPPORTUNISTIC MACROALGAE

Opportunistic macroalgae are a primary symptom of estuary eutrophication. They are highly effective at utilising excess nitrogen, enabling them to out-compete other seaweed species and, at nuisance levels, can form mats on the estuary surface which adversely impact underlying sediments and fauna, other algae, fish, birds, seagrass, and saltmarsh. Macroalgae that becomes detached can also accumulate and decay in subtidal areas and on shorelines causing oxygen depletion and nuisance odours and conditions. The greater the density, persistence, and extent of macroalgal entrainment within sediments, the greater the subsequent impacts.

Opportunistic macroalgal growth was assessed by mapping the spatial spread and density of macroalgae in the Available Intertidal Habitat (AIH) (Figure 7), and calculating an "Ecological Quality Rating" (EQR) using the Opportunistic Macroalgal Blooming Tool (OMBT).

The EQR score can range from zero (major disturbance) to one (reference/minimally disturbed) and relates to a quality status threshold band (i.e. bad, poor, good, moderate, high - Section 2, Table 2, Appendix 4). The individual metrics that are used to calculate the EQR (spatial extent, density, biomass, and degree of sediment entrainment of macroalgae within the affected intertidal area), are also scored and have quality status threshold bands to guide key drivers of change. If the estuary supports <5% opportunistic macroalgal cover within the AIH, the overall quality status is reported as "HIGH" with no further sampling required.



4. RESULTS AND DISCUSSION (CONTINUED)




Figure 6. Map of areas with low sediment oxygenation - Waikouaiti Estuary, Dec. 2016.

4. RESULTS AND DISCUSSION (CONTINUED)

The overall opportunistic macroalgal EQR for Waikouaiti Estuary in Dec. 2016 was 0.69 (Table 6), a quality status of “GOOD” and indicates that the estuary overall is expressing limited symptoms of eutrophication. This is apparent through the “Good” Quality Status scores for the affected area of the estuary, the relatively low overall percentage cover and biomass in available intertidal habitat, and the absence of sediment entrained algae (“High” Quality Status). However, there are indications of excessive growth present with high biomass in affected areas. In other words, macroalgal growth is not widespread, but when it is present, it can be present in high densities (e.g. Figure 7).

Growths were dominated by the green alga *Ulva intestinalis* with the red alga *Gracilaria chilensis* an occasional subdominant. The highest density growths were located on the strongly flushed intertidal flats of the main estuary channel. The regular flushing of these flats with waters containing elevated nutrients appears sufficient to stimulate growth, but the extent that macroalgae can establish nuisance beds is limited by the absence of hard substrate to adhere to and regular flushing limiting the buildup of fine muds that often result in sediment entrained growths of *Gracilaria* which can lead to significant sediment anoxia and enrichment.

While extensive dense beds were present in some areas, and were also evident as prolific subtidal growths, the combined overall risk rating was assessed as LOW at the time of sampling. It is noted however that the estuary has previously shown excessive macroalgal growth including nuisance red algal blooms in some years. Data collected by the authors in 2013 showed dense blooms of the red opportunistic macroalgae *Gracilaria chilensis*, with a subdominant cover of *Ulva intestinalis* (>95% cover and mean biomass of 3500gm⁻² wet wgt.), with “gross eutrophic zone” conditions present e.g. anoxic sediments, soft muds and high nuisance macroalgal cover. Previous monitoring in 2006 reported 0.6ha of macroalgae in the estuary (Stewart 2007).




Table 6. Summary of intertidal opportunistic macroalgal cover, Waikouaiti Estuary, Dec. 2016.

Metric	Face Value	Final Equidistant Score (FEDS)	Quality Status
AIH - Available Intertidal Habitat (ha)	97		
Percentage cover of AIH (%) = (Total % Cover / AIH} x 100 <i>where Total % cover = Sum of {(patch size) / 100} x average % cover for patch</i>	7.8	0.74	Good
Biomass of AIH (g.m ⁻²) = Total biomass / AIH <i>where Total biomass = Sum of (patch size x average patch biomass)</i>	118.0	0.76	Good
Biomass of Affected Area (g.m ⁻²) = Total biomass / AA <i>where Total biomass = Sum of (>5% cover patch size x average patch biomass)</i>	524.2	0.39	Poor
Presence of Entrained Algae = (No. quadrats or area (ha) with entrained algae / total no. of quadrats or area (ha)) x 100	0.0	1.00	High
Affected Area (use the lowest of the following two metrics)		0.56	Moderate
Affected Area, AA (ha) = Sum of all patch sizes (with macroalgal cover >5%)	21.8	0.74	Good
Size of AA in relation to AIH (%) = (AA / AIH) x 100	22.5	0.56	Moderate
OVERALL MACROALGAL ECOLOGICAL QUALITY RATING - EQR (AVERAGE OF FEDS)		0.69	GOOD
Upper and Lower 90% Confidence Interval for EQR		0.633	0.816

4. RESULTS AND DISCUSSION (CONTINUED)




Figure 7. Map of intertidal opportunistic macroalgal biomass (g.m⁻²) - Waikouaiti Estuary, Dec. 2016.

4. RESULTS AND DISCUSSION (CONTINUED)

4.5. SEAGRASS



Seagrass (*Zostera muelleri*) beds are important ecologically because they enhance primary production and nutrient cycling, stabilise sediments, elevate biodiversity, and provide nursery and feeding grounds for a range of invertebrates and fish. Though tolerant of a wide range of conditions, seagrass is vulnerable to excessive nutrients, fine sediments in the water column, and sediment quality (particularly if there is a lack of oxygen and production of sulphides).

Table 7 and Figure 8 summarise the results of the intertidal seagrass extent (the mapped intertidal estuary area minus saltmarsh). The results show:

- 2% of the intertidal estuary area (1.7ha) supported seagrass growth (0.2ha was recorded in 2006).
- All seagrass beds were located in muddy sand substrate in the main river channel of the central estuary.
- Seagrass was present in two density categories, dense beds (80% cover), and moderately dense beds (50% cover.)

The extent of intertidal seagrass in Waikouaiti Estuary is relatively small and confined to areas near the main channel in the central basin that have strong tidal flushing and are subsequently not prone to excessive fine mud deposition. However, the seagrass beds were clearly under stress from smothering macroalgal growths in many places. The absence of seagrass within the Merton and East arms is considered likely due to influence of elevated mud in these areas combined with relatively long tidal exposure periods due to their relatively perched nature creating conditions unsuitable for seagrass growth e.g. extended exposure periods resulting in plant stress from dessication.

In the absence of any comprehensive rating of seagrass extent within NZ estuaries, which can be highly variable in the extent of seagrass that they support, changes from a documented baseline currently represent the most reliable method for monitoring seagrass extent and assessing change. The current study has provided a high resolution GIS map of seagrass extent for this purpose.

Table 7. Summary of seagrass (*Z. muelleri*) cover, Waikouaiti Estuary, 2016.

Percentage Cover	Area (ha)	Percentage
0	94.9	98.2
50%	1.6	1.6
80%	0.1	0.1
	96.6	100



4. RESULTS AND DISCUSSION (CONTINUED)




Figure 8. Map of intertidal seagrass percentage cover (*Zostera muelleri*) - Waikouaiti Estuary, 2016.

4. RESULTS AND DISCUSSION (CONTINUED)

4.6. SALTMARSH



Saltmarsh (vegetation able to tolerate saline conditions where terrestrial plants are unable to survive) is important as it is highly productive, naturally filters and assimilates sediment and nutrients, acts as a buffer that protects against introduced grasses and weeds, and provides an important habitat for a variety of species including fish and birds. Saltmarsh generally has the most dense cover in the sheltered and more strongly freshwater influenced upper estuary, and relatively sparse cover in the lower (more exposed and saltwater dominated) parts of the estuary, with the lower extent of saltmarsh growth limited for most species to above the height of mean high water neap (MHWN).

The primary measure to assess saltmarsh condition is the percent cover of the intertidal area. Table 8 and Figure 9 summarise the 2016 results and show saltmarsh was present across 80.3ha (45%) of the intertidal estuary area, a risk indicator rating of LOW. Saltmarsh was dominated by herbfields (96%) located in extensive beds in the sheltered arms. Glasswort was the dominant cover, commonly mixed with primrose, and remuremu, with introduced weeds and grasses common in the upper tidal range, particularly where drainage channels have been modified. There were only small areas dominated by rushland (jointed wirerush - 2.4% predominantly in the Merton arm) or reedland (the introduced invasive cord grass *Spartina* - 1.4% predominantly in the East arm), (Figure 7). Elsewhere channelling of the main river, or steep edges to reclaimed or drained estuary margins, restricted saltmarsh to a relatively narrow strip along the upper tidal reaches (see sidebar photos).

A supporting measure also applied is saltmarsh loss compared to estimated natural state cover. While the historical extent of the estuary has not been specifically mapped as part of the current work, it is evident that extensive areas in the upper estuary have been historically drained and converted to pasture. It is estimated that ~150ha of saltmarsh has been lost from the estuary, of this ~31ha in the past decade, a supporting risk rating of HIGH. The combined overall risk rating was assessed as LOW to MODERATE recognising that while historical losses have been significant, and appear to be ongoing, saltmarsh remains a significant feature of the estuary.

Table 8. Summary of dominant saltmarsh cover, Waikouaiti Estuary, 2016.

Class	Dominant Species	Primary subdominant species	Area (ha)	Percentage
Reedland			1.2	1.4%
	<i>Spartina sp. (Cord grass)</i>		1.2	
Rushland			1.9	2.4%
	<i>Apodasmia similis (Jointed wirerush)</i>		1.9	
Herbfield			77.2	96.1%
	<i>Sarcocornia quinqueflora (Glasswort)</i>	<i>Sarcocornia quinqueflora (Glasswort)</i>	39.8	
		<i>Samolus repens (Primrose)</i>	4.9	
		<i>Selliera radicans (Remuremu)</i>	20.1	
		<i>Introduced weeds</i>	12.4	
Total (Ha)			80.3	

4. RESULTS AND DISCUSSION (CONTINUED)


- Waikouaiti 2017**
Saltmarsh Class
-  Estuarine Shrub
 -  Tussockland
 -  Sedgeland
 -  Reedland
 -  Rushland
 -  Herbfield



Figure 9. Map of dominant saltmarsh cover - Waikouaiti Estuary, Dec. 2016.

4. RESULTS AND DISCUSSION (CONTINUED)

4.7. 200m TERRESTRIAL MARGIN



Like saltmarsh, a densely vegetated terrestrial margin filters and assimilates sediment and nutrients, acts as an important buffer that protects against introduced grasses and weeds, is an important habitat for a variety of species, provides shade to help moderate stream temperature fluctuations, and improves estuary biodiversity. The results of the 200m terrestrial margin mapping of the estuary (Table 9 and Figure 10) showed:

- 75% was pasture or unmaintained grassland.
- 13% had been developed (residential, industrial, road and rail).
- 12% was densely vegetated.

A similar pattern is evident on a catchment wide scale with 74% of the catchment developed into high producing grassland (Figure 11).

The 200m terrestrial margin estuary margin itself has been significantly modified historically, primarily through reclamation and drainage for conversion to pasture, armouring for flood control and road and rail infrastructure, and residential development.

The small remaining extent of densely vegetated 200m terrestrial margin habitat means there is very limited buffering against adverse ecological degradation (e.g. localised sediment and nutrient inputs, introduced weeds), and poor natural ecological connectivity between the estuary and surrounding natural habitats. The 200m terrestrial margin risk indicator rating is therefore "HIGH".

Table 9. Summary of 200m terrestrial margin land cover, Waikouaiti Estuary, 2016.

Class	Dominant features	Percentage
Forest	Mixed native and exotic trees predominantly on the coastal spit and in narrow margin strips near the estuary margin. Common species include <i>Pinus radiata</i> (Pine tree), <i>Ulex europaeus</i> (Gorse), <i>Lupinus arboreus</i> (Tree lupin), <i>Ammophila arenaria</i> (Marram grass)	0.2
Scrub/Forest		6.1
Scrub		5.6
Duneland	Small area of marram grass on the coastal spits near the entrance	0.1
Pasture	Developed pasture and un-maintained grassland present along much of the estuary edge	47.5
Grassland		27.1
Industrial	Sewage ponds, commercial chicken farming	1.4
Residential	Karitane	9.3
Built feature	Road and rail corridors, wharf	2.7
Total		100



Grassland bordering estuary margins



Developed pasture to estuary margins. Modified estuary drainage also evident.



Grassland along road edge at estuary margin.

4. RESULTS AND DISCUSSION (CONTINUED)



Figure 10. Map of 200m Terrestrial Margin - Dominant Land Cover, Waikouaiti Estuary, Dec. 2016.

4. RESULTS AND DISCUSSION (CONTINUED)




Figure 11. Summary of Catchment Land Cover (LCDB4 2012), Waikouaiti Estuary.

4. RESULTS AND DISCUSSION (CONTINUED)

4.8. NZ ESTUARY TROPHIC INDEX

The recently developed NZ ETI (Robertson et al. 2016a,b) is designed to enable the consistent assessment of estuary state in relation to nutrient enrichment, and also includes assessment criteria for sediment muddiness. A key part of the ETI output has been the development of an integrated calculator that enables easy calculation of estuary physical and nutrient load susceptibility (primarily based on catchment nutrient loads combined with mixing and dilution in the estuary), as well as trophic expression based on key estuary indicators.

From the current broad scale monitoring (this report) and fine scale monitoring (Robertson et al. 2017), Table 10 presents the indicators used to derive an ETI score for the estuary.

The physical and nutrient load susceptibility of Waikouaiti Estuary has been determined as "HIGH" based on catchment estimates of nutrient loads derived from NIWAs CLUES model, estuary characteristics sourced from the Coastal Explorer database, and ecological value. The overall ETI score for the estuary is 0.51, a risk rating of "MODERATE" for eutrophic symptoms.

Table 10. Primary and supporting indicator values used to calculate an ETI score for Waikouaiti Estuary, December 2016.

PRIMARY SYMPTOM INDICATORS FOR SHALLOW INTERTIDAL DOMINATED ESTUARIES (AT LEAST 1 PRIMARY SYMPTOM INDICATOR REQUIRED)				Primary Symptom Value
Required	Opportunistic Macroalgae	OMBT EQR	shallow inter-tidal	0.69
	Macroalgal GEZ/Estuary Area	% Gross Eutrophic Zone (GEZ)		0
	Macroalgal GEZ	Ha Gross Eutrophic Zone (GEZ)		0
Optional	Phytoplankton biomass	Chl- a (summer 90 pct, mg/m ³)	water column	<1*
	Cyanobacteria (if issue identified)			not yet developed
SUPPORTING INDICATORS FOR SHALLOW INTERTIDAL DOMINATED ESTUARIES (MUST INCLUDE A MINIMUM OF 1 REQUIRED INDICATOR)				Supporting Indicator Value
Required Indicators	Sediment Oxygenation	Mean Redox Potential (mV) at 1cm depth in most impacted sediments and representing at least 10% of estuary area	shallow inter-tidal	-359mV
		% of estuary with Redox Potential <-150mV at 3cm		10%
		Ha of estuary with Redox Potential <-150mV at 3cm		24
		% of estuary with apparent Redox Potential Depth <1cm		-
	Sediment Total Organic Carbon	Mean TOC (%) measured at 0-2cm depth in most impacted sediments and representing at least 10% of estuary area		0.4
	Sediment Total Nitrogen	Mean TN (mg/kg) measured at 0-2cm depth in most impacted sediments and representing at least 10% of estuary area		633
	Macroinvertebrates	Mean AMBI score measured at 0-15cm depth in most impacted sediments and representing at least 10% of estuary area		4.5
		% of estuary with AMBI >4.3		not yet developed
		TBI (if toxicity an issue)		not yet developed
	Optional Indicators	Dissolved oxygen		7 day mean, 7 day mean minimum, 1 day minimum (mg/m ³)
TN and TP concentration		mg/l - 7 day mean, 7 day mean minimum, 1 day minimum	not yet developed	
Water Clarity		Secchi or black disc (m)	not yet developed	
Sediment Sulphur		Requires development	shallow inter-tidal	not yet developed
Soft mud		% estuary area with soft mud (>25% mud content)		21%
		% mud content (mean of whole estuary area)		not yet developed
Sedimentation Rate		Ratio of current annual mean relative to Natural Sed. Rate (NSR)		4.6
		% Estuary Area with Sedimentation Rate >5xNSR	baseline estab.	
SAV (Seagrass)	Extent (% of ENSC)	all habitat	-	
	% change from measured baseline		baseline estab.	
NZ ETI Score				0.51

*surface and bottom measurements at 3 sites collected on 8/12/17 (Robertson et al. 2017)

5. SUMMARY AND CONCLUSIONS

Broad scale habitat mapping undertaken in December 2016, combined with ecological risk indicator ratings in relation to the key estuary stressors (i.e. muddiness, eutrophication and habitat modification) have been used to assess overall estuary condition.

Muddiness

Soft or very soft muds covered 20.9ha (21.7%) of the intertidal area, a risk indicator rating of HIGH, and had a mud content measured in representative areas of 20-29%, a supporting risk indicator rating of MODERATE to HIGH. 24ha (10%) of the intertidal area (including saltmarsh) had sediment oxygenation depleted to a level where adverse impacts to macrofauna (sediment and surface dwelling animals) are expected, an ETI risk indicator rating of HIGH. Soft muds were concentrated in the upper tidal reaches of the estuary embayments where mud settlement is thought to predominantly reflect the presence of sheltered deposition zones in the Merton and East arms, to a lesser extent, salinity driven flocculation. The strongly flushed main river channel habitat in relatively good condition with limited accumulation of muds and generally good sediment oxygenation.

Eutrophication

The NZ ETI combines a range of broad and fine scale indicators (see also Robertson et al. 2017) to provide an overall assessment of eutrophic expression in the estuary, including primary productivity through macroalgal growth and phytoplankton, and supporting indicators of sediment muddiness, oxygenation, organic content, nutrients, macroinvertebrates, and the presence of gross eutrophic zones (a combined presence of dense macroalgal growth, muds and poor sediment oxygenation). The overall ETI score for the estuary in December 2016 was 0.51, a risk rating of MODERATE for eutrophic symptoms.

Although nuisance macroalgal growths had not resulted in the establishment of gross eutrophic conditions in 2016, there were prolific *Ulva* growths in parts of the main channel, and in 2013 nuisance *Gracilaria* growths and gross eutrophic zones were present in the Merton and East arms. Such findings highlight that nutrient inputs to the estuary are sufficient to fuel nuisance growths and that at times the estuary shifts to a more degraded state. For example the NZ ETI score for 2013 was 0.90, a risk rating of HIGH for eutrophic symptoms.

Habitat modification

Despite significant historical losses of saltmarsh (estimated at ~150ha in total with ~30 in the past 10 years), extensive herbfield-dominated saltmarsh remained in the estuary (80ha, 45% of the intertidal area). The 200m terrestrial margin had also been highly modified with 75% pasture or unmaintained grassland, 13% developed (residential/road/rail), and only 12% supported a densely vegetated buffer.

The estuary supported small areas of high value seagrass beds (1.7ha, 2%) near the main channel in the central basin where there is strong tidal flushing and little fine mud deposition. However, the seagrass beds were clearly under stress from smothering macroalgal growths in many places.

Comparison with 2006 results

A comparison of the 2006 and 2016 results show potential improvements in the estuary from a decrease in the area of soft mud (39ha in 2006 to 21ha in 2016) and an increase in seagrass (0.2ha to 1.7ha), but declines in quality related to increased macroalgal cover (0.6 ha to 5.2ha), and a significant decrease in saltmarsh (111ha to 80ha). Future monitoring will determine if these results reflect ongoing trends in broad scale features of the estuary.

The combined results place the estuary in a "MODERATE" state overall in relation to ecological health. Fine sediment issues are evident in the relatively sheltered Merton and East arms, and eutrophication issues have been previously apparent in these same locations. Dense opportunistic macroalgal growths were apparent in the main estuary channel. The modification and loss of estuary saltmarsh and a densely vegetated buffer have been historically significant, and ongoing pressures are evident in these areas.

6. MONITORING RECOMMENDATIONS

Waikouaiti Estuary has been identified by ORC as a priority for monitoring because it is a moderate-large estuary with high ecological and human use values that is situated in a developed catchment, and therefore vulnerable to excessive sedimentation and eutrophication. As a consequence, it is a key part of ORC's coastal monitoring programme being undertaken throughout the Otago region. Broad scale habitat mapping and the first year of baseline fine scale sampling have now been undertaken (December 2016).

In order to assess ongoing long-term trends in the condition of such estuaries, it is common practice amongst NZ Regional Councils to establish a strong baseline against which future trends can be compared. This typically comprises comprehensive broad scale habitat mapping on a 5-10 yearly cycle, targeted annual monitoring where specific issues are identified (e.g. opportunistic nuisance macroalgal growth), and fine scale monitoring comprising 3-4 consecutive years of baseline monitoring, followed by 5 yearly impact monitoring.

The present report addresses the broad scale component of the long term programme. The recommendation for ongoing monitoring to meet this requirement for the Waikouaiti Estuary is as follows:

Broad Scale Habitat Mapping

To characterise any issues of change in habitat (e.g. saltmarsh or seagrass area, soft mud extent), it is recommended that broad scale habitat mapping be undertaken at 10 yearly intervals (next scheduled for 2026) unless obvious changes are observed in the interim.

To fully characterise the observed variation between years in opportunistic macroalgal growth (both green and red blooms), it is recommended that a strong baseline be established by undertaking an annual macroalgal survey (January/February) over the next three years in conjunction with scheduled fine scale sampling.

Fine Scale Monitoring

To complete the fine scale baseline in Waikouaiti Estuary, it is recommended that the remaining 3 consecutive years of annual summer (i.e. December-February) fine scale monitoring of intertidal sites (including sedimentation rate measures), be undertaken in 2017, 2018 and 2019 (preferably during a summer low flow period).

To fully characterise the potential for upper estuary stratification and eutrophication, it is recommended that water column monitoring of the upper to mid estuary be undertaken during a prolonged summer, low flow period in 2018. It is envisaged that this should include sampling of surface and bottom water at 5-6 sites, between the Railway Bridge and 1km below Orbells Crossing.

To characterise the potential for excessive sedimentation, it is recommended that sedimentation rate be assessed annually, using appropriately placed sediment plates, and areal extent of muddy sediments be assessed at 5-10 yearly intervals (the latter assessed in broad scale monitoring).



7. ACKNOWLEDGEMENTS

This survey and report has been undertaken with the support and assistance of Rachel Ozanne (Coastal Scientist, ORC). Her review of this report was much appreciated. We are also very grateful to Ben Robertson and Maz Robertson (Wriggle) for help with the field sampling.

8. REFERENCES

- Atkinson, I.A.E. 1985. Derivation of vegetation mapping units for an ecological survey of Tongariro National Park Nth Island, NZ. *NZ Journal of Botany*, 23; 361-378.
- Birchenough, S., Parker N., McManus E. and Barry, J. 2012. Combining bioturbation and redox metrics: potential tools for assessing seabed function. *Ecological Indicators* 12: 8-16.
- Davey, A. 2009. Confidence of Class for WFD Marine Plant Tools. WRC report EA7954. 34pp.
- Ellis, J., Cummings, V., Hewitt, J., Thrush, S., Norkko, A. 2002. Determining effects of suspended sediment on condition of a suspension feeding bivalve (*Atrina zelandica*): results of a survey, a laboratory experiment and a field transplant experiment. *Journal of Experimental Marine Biology and Ecology*, 267, 147–174.
- Fenchel, T. and Riedl, R. 1970. The sulphide system: a new biotic community underneath the oxidized layer of marine sand bottoms. *Mar Biol* 7: 255-268.
- Hargrave, B.T., Holmer, M. and Newcombe, C.P. 2008. Towards a classification of organic enrichment in marine sediments based on biogeochemical indicators. *Marine Pollution Bulletin*, 56(5), pp.810–824.
- Hunting, E.R. and Kampfraath, A.A. 2012. Contribution of bacteria to redox potential (E_h) measurements in sediments. *International Journal of Environmental Science and Technology*, 10(1): 55-62.
- Jørgensen, N. and Revsbech, N.P. 1985. Diffusive boundary layers and the oxygen uptake of sediments and detritus. *Limnology and Oceanography* 30:111-122.
- Keeley, N.B. et al. 2012. Exploiting salmon farm benthic enrichment gradients to evaluate the regional performance of biotic indices and environmental indicators. *Ecological Indicators*, 23, pp.453–466.
- Lohrer, A., Thrush, S., Hewitt, J., Berkenbusch, K., Ahrens, M., Cummings, V. 2004. Terrestrially derived sediment: response of marine macrobenthic communities to thin terrigenous deposits. *Marine Ecology Progress Series*, 273, 121–138.
- Mannino, A. and Montagna, P. 1997. Small-Scale Spatial Variation of Macrobenthic Community. *Estuaries*, 20, 159–173.
- Nelson, Walter G. (ed.) 2009. *Seagrasses and Protective Criteria: A Review and Assessment of Research Status*. Office of Research and Development, National Health and Environmental Effects Research Laboratory, EPA/600/R-09/050.
- Norkko, A., Talman, S., Ellis, J., Nicholls, P. and Thrush, S. 2002. Macrofaunal Sensitivity to Fine Sediments in the Whitford Embayment. Auckland Regional Council, Technical Publication, 158, 1–30.
- Peeters, E., Gardeniers, J., Koelmans, A. 2000. Contribution of trace metals in structuring in situ macroinvertebrate community composition along a salinity gradient. *Environmental Toxicology and Chemistry*, 19, 1002–1010.
- Rakocinski, C., Brown, S., Gaston, G., Heard, R., Walker, W. and Summers, J. 1997. Macrobenthic Responses to Natural and Contaminant-Related Gradients in Northern Gulf of Mexico Estuaries. *Ecological Applications*, 7, 1278–1298.
- Revsbech, N.P., Sørensen, J., Blackburn, T.H. and Lomholt, J.P. 1980. Distribution of oxygen in marine sediments measured with microelectrodes. *Limnology and Oceanography* 25: 403-411.
- Robertson, B.M., Gillespie, P.A., Asher, R.A., Frisk, S., Keeley, N.B., Hopkins, G.A., Thompson, S.J., Tuckey, B.J. 2002. *Estuarine Environmental Assessment and Monitoring: A National Protocol. Part A. Development, Part B. Appendices, and Part C. Application*. Prepared for supporting Councils and the Ministry for the Environment, Sustainable Management Fund Contract No. 5096. Part A. 93p. Part B. 159p. Part C. 40p plus field sheets.
- Robertson, B.M., Robertson, B.P., and Stevens, L.M. 2017. Waikouaiti Estuary: Fine Scale Monitoring 2016/17. Report prepared by Wriggle Coastal Management for Otago Regional Council. 37p.
- Robertson, B.M., Stevens, L., Robertson, B.P., Zeldis, J., Green, M., Madarasz-Smith, A., Plew, D., Storey, R., Hume, T. and Oliver, M. 2016a. NZ Estuary Trophic Index. Screening Tool 1. Determining eutrophication susceptibility using physical and nutrient load data. Prepared for Envirolink Tools Project: Estuarine Trophic Index MBIE/NiWA Contract No: C01X1420. 47p.
- Robertson, B.M., Stevens, L., Robertson, B.P., Zeldis, J., Green, M., Madarasz-Smith, A., Plew, D., Storey, R., Hume, T. and Oliver, M. 2016b. NZ Estuary Trophic Index. Screening Tool 2. Screening Tool 2. Determining Monitoring Indicators and Assessing Estuary Trophic State. Prepared for Envirolink Tools Project: Estuarine Trophic Index MBIE/NiWA Contract No: C01X1420. 68p.
- Robertson, B.P., Gardner, J.P.A., Savage, C., Robertson, B.M. and Stevens, L.M. 2016. Optimising a widely-used coastal health index through quantitative ecological group classifications and associated thresholds. *Ecological Indicators*. 69. 595-605.
- Robertson, B.P., Gardner, J.P.A. and Savage, C. 2015. Macrobenthic - mud relations strengthen the foundation for benthic index development : A case study from shallow, temperate New Zealand estuaries. *Ecological Indicators*, 58, pp.161–174. Available at: <http://dx.doi.org/10.1016/j.ecolind.2015.05.039>.
- Robertson, B.P. 2013. Determining the sensitivity of macroinvertebrates to fine sediments in representative New Zealand estuaries. Honours dissertation, Victoria University of Wellington.

9. REFERENCES (CONTINUED)

- Rosenberg, R., Nilsson, H.C. and Diaz, R.J. 2001. Response of benthic fauna and changing sediment redox profiles over a hypoxic gradient. *Estuarine Coast Shelf Science* 53: 343-350.
- Sakamaki, T., Nishimura, O. 2009. Is sediment mud content a significant predictor of macrobenthos abundance in low-mud-content tidal flats? *Marine and Freshwater Research*, 60, 160.
- Thrush, S.F., Hewitt, J., Norkko, A., Nicholls, P., Funnell, G. and Ellis, J. 2003. Habitat change in estuaries: predicting broad-scale responses of intertidal macrofauna to sediment mud content. *Marine Ecology Progress Series* 263, 101–112.
- Wehkamp, S., Fischer, P. 2012. Impact of hard-bottom substrata on the small-scale distribution of fish and decapods in shallow subtidal temperate waters. *Helgoland Marine Research*, 67, 59–72.
- WFD-UKTAG (Water Framework Directive – United Kingdom Technical Advisory Group). (2014). UKTAG Transitional and Coastal Water Assessment Method Macroalgae Opportunistic Macroalgal Blooming Tool. Retrieved from [http://www.wfd.uk.org/sites/default/files/Media/Characterisation of the water environment/Biological Method Statements/TraC Macroalgae OMBT UKTAG Method Statement.PDF](http://www.wfd.uk.org/sites/default/files/Media/Characterisation%20of%20the%20water%20environment/Biological%20Method%20Statements/TraC%20Macroalgae%20OMBT%20UKTAG%20Method%20Statement.PDF).

References for Table 1

- Abraham, G. 2005. *Holocene sediments of Tamaki Estuary: characterisation and impact of recent human activity on an urban estuary in Auckland, NZ*. PhD Thesis, University of Auckland, Auckland, NZ, p 361.
- Anderson, D., Gilbert, P. and Burkholder, J. 2002. Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries* 25, 704–726.
- Ferreira, J., Andersen, J. and Borja, A. 2011. Overview of eutrophication indicators to assess environmental status within the European Marine Strategy Framework Directive. *Estuarine, Coastal and Shelf Science* 93, 117–131.
- Gibb, J.G. and Cox, G.J. 2009. *Patterns & Rates of Sedimentation within Porirua Harbour*. Consultancy Report (CR 2009/1) prepared for Porirua City Council. 38p plus appendices.
- IPCC. 2007. Intergovernmental Panel on Climate Change web site. https://www.ipcc.ch/publications_and_data/ar4/wg1/ (accessed December 2009).
- IPCC. 2013. Intergovernmental Panel on Climate Change web site. <https://www.ipcc.ch/report/ar5/wg1/> (accessed March 2014).
- Kennish, M.J. 2002. *Environmental threats and environmental future of estuaries*. *Environmental Conservation* 29, 78–107.
- National Research Council. 2000. *Clean coastal waters: understanding and reducing the effects of nutrient pollution*. Ocean Studies Board and Water Science and Technology Board, Commission on Geosciences, Environment, and Resources. Washington, DC: National Academy Press. 405p.
- Painting, S.J., Devlin, M.J., Malcolm, S.J., Parker, E.R., Mills, D.K., Mills, C., and Winpenny, K. 2007. Assessing the impact of nutrient enrichment in estuaries: susceptibility to eutrophication. *Marine Pollution Bulletin* 55(1-6), 74–90.
- Robertson, B.M. and Stevens, L.M. 2007. *Waikawa Estuary 2007 Fine Scale Monitoring and Historical Sediment Coring*. Prepared for Environment Southland. 29p.
- Robertson, B.M. and Stevens, L.M. 2010. *New River Estuary: Fine Scale Monitoring 2009/10*. Report prepared by Wriggle Coastal Management for Environment Southland. 35p.
- de Salas, M.F., Rhodes, L.L., Mackenzie, L.A., Adamson, J.E. 2005. Gymnodinoid genera *Karenia* and *Takayama* (Dinophyceae) in New Zealand coastal waters. *New Zealand Journal of Marine and Freshwater Research* 39, 135–139.
- Stewart, J.R., Gast, R.J., Fujioka, R.S., Solo-Gabriele, H.M., Meschke, J.S., Amaral-Zettler, L.A., Castillo, E. Del., Polz, M.F., Collier, T.K., Strom, M.S., Sinigalliano, C.D., Moeller, P.D.R. and Holland, A.F. 2008. The coastal environment and human health: microbial indicators, pathogens, sentinels and reservoirs. *Environmental Health* 7 Suppl 2, S3.
- Swales, A., and Hume, T. 1995. *Sedimentation history and potential future impacts of production forestry on the Wharekawa Estuary, Coromandel Peninsula*. Prepared for Carter Holt Harvey Forests Ltd. NIWA report no. CHH004.
- Valiela, I., McClelland, J., Hauxwell, J., Behr, P., Hersh, D., and Foreman, K. 1997. Macroalgal blooms in shallow estuaries: Controls and ecophysiological and ecosystem consequences. *Limnology and Oceanography* 42, 1105–1118.
- Wade, T.J., Pai, N., Eisenberg, J.N.S., and Colford, J.M., 2003. Do U.S. Environmental Protection Agency Water Quality Guidelines for Recreational Waters Prevent Gastrointestinal Illness? A Systematic Review and Meta-analysis. *Environmental Health Perspective* 111, 1102–1109.

APPENDIX 1. BROAD SCALE HABITAT CLASSIFICATION DEFINITIONS.

Vegetation was classified using an interpretation of the Atkinson (1985) system, whereby dominant plant species were coded by using the two first letters of their Latin genus and species names e.g. marram grass, *Ammophila arenaria*, was coded as Amar. An indication of dominance is provided by the use of () to distinguish subdominant species e.g. Amar(Caed) indicates that marram grass was dominant over ice plant (*Carpobrotus edulis*). The use of () is not always based on percentage cover, but the subjective observation of which vegetation is the dominant or subdominant species within the patch. A measure of vegetation height can be derived from its structural class (e.g. rushland, scrub, forest).

- Forest:** Woody vegetation in which the cover of trees and shrubs in the canopy is >80% and in which tree cover exceeds that of shrubs. Trees are woody plants ≥ 10 cm diameter at breast height (dbh). Tree ferns ≥ 10 cm dbh are treated as trees. Commonly sub-grouped into native, exotic or mixed forest.
- Treeland:** Cover of trees in the canopy is 20–80%. Trees are woody plants > 10 cm dbh. Commonly sub-grouped into native, exotic or mixed treeland.
- Scrub:** Cover of shrubs and trees in the canopy is >80% and in which shrub cover exceeds that of trees (c.f. FOREST). Shrubs are woody plants < 10 cm dbh. Commonly sub-grouped into native, exotic or mixed scrub.
- Shrubland:** Cover of shrubs in the canopy is 20–80%. Shrubs are woody plants < 10 cm dbh. Commonly sub-grouped into native, exotic or mixed shrubland.
- Tussockland:** Vegetation in which the cover of tussock in the canopy is 20–100% and in which the tussock cover exceeds that of any other growth form or bare ground. Tussock includes all grasses, sedges, rushes, and other herbaceous plants with linear leaves (or linear non-woody stems) that are densely clumped and > 100 cm height. Examples of the growth form occur in all species of *Cortaderia*, *Gahnia*, and *Phormium*, and in some species of *Chionochloa*, *Poa*, *Festuca*, *Rytidosperma*, *Cyperus*, *Carex*, *Uncinia*, *Juncus*, *Astelia*, *Aciphylla*, and *Celmisia*.
- Duneland:** Vegetated sand dunes in which the cover of vegetation in the canopy (commonly Spinifex, Pingao or Marram grass) is 20–100% and in which the vegetation cover exceeds that of any other growth form or bare ground.
- Grassland:** Vegetation in which the cover of grass (excluding tussock-grasses) in the canopy is 20–100%, and in which the grass cover exceeds that of any other growth form or bare ground.
- Sedgeland:** Vegetation in which the cover of sedges (excluding tussock-sedges and reed-forming sedges) in the canopy is 20–100% and in which the sedge cover exceeds that of any other growth form or bare ground. "Sedges have edges." Sedges vary from grass by feeling the stem. If the stem is flat or rounded, it's probably a grass or a reed, if the stem is clearly triangular, it's a sedge. Sedges include many species of *Carex*, *Uncinia*, and *Scirpus*.
- Rushland:** Vegetation in which the cover of rushes (excluding tussock-rushes) in the canopy is 20–100% and where rush cover exceeds that of any other growth form or bare ground. A tall grasslike, often hollow-stemmed plant, included in rushland are some species of *Juncus* and all species of *Leptocarpus*.
- Reedland:** Vegetation in which the cover of reeds in the canopy is 20–100% and in which the reed cover exceeds that of any other growth form or open water. Reeds are herbaceous plants growing in standing or slowly-running water that have tall, slender, erect, unbranched leaves or culms that are either round and hollow – somewhat like a soda straw, or have a very spongy pith. Unlike grasses or sedges, reed flowers will each bear six tiny petal-like structures. Examples include *Typha*, *Bolboschoenus*, *Scirpus lacustris*, *Eleocharis sphacelata*, and *Baumea articulata*.
- Cushionfield:** Vegetation in which the cover of cushion plants in the canopy is 20–100% and in which the cushion-plant cover exceeds that of any other growth form or bare ground. Cushion plants include herbaceous, semi-woody and woody plants with short densely packed branches and closely spaced leaves that together form dense hemispherical cushions.
- Herbfield:** Vegetation in which the cover of herbs in the canopy is 20–100% and where herb cover exceeds that of any other growth form or bare ground. Herbs include all herbaceous and low-growing semi-woody plants that are not separated as ferns, tussocks, grasses, sedges, rushes, reeds, cushion plants, mosses or lichens.
- Lichenfield:** Vegetation in which the cover of lichens in the canopy is 20–100% and where lichen cover exceeds that of any other growth form or bare ground.
- Introduced weeds:** Vegetation in which the cover of introduced weeds in the canopy is 20–100% and in which the weed cover exceeds that of any other growth form or bare ground.
- Seagrass meadows:** Seagrasses are the sole marine representatives of the Angiospermae. They all belong to the order Helobiae, in two families: Potamogetonaceae and Hydrocharitaceae. Although they may occasionally be exposed to the air, they are predominantly submerged, and their flowers are usually pollinated underwater. A notable feature of all seagrass plants is the extensive underground root/rhizome system which anchors them to their substrate. Seagrasses are commonly found in shallow coastal marine locations, salt-marshes and estuaries and is mapped separately to the substrates they overlie.
- Macroalgal bed:** Algae are relatively simple plants that live in freshwater or saltwater environments. In the marine environment, they are often called seaweeds. Although they contain chlorophyll, they differ from many other plants by their lack of vascular tissues (roots, stems, and leaves). Many familiar algae fall into three major divisions: Chlorophyta (green algae), Rhodophyta (red algae), and Phaeophyta (brown algae). Macroalgae are algae observable without using a microscope. Macroalgal density, biomass and entrainment are classified and mapped separately to the substrates they overlie.
- Cliff:** A steep face of land which exceeds the area covered by any one class of plant growth-form. Cliffs are named from the dominant substrate type when unvegetated or the leading plant species when plant cover is $\geq 1\%$.
- Rock field:** Land in which the area of residual rock exceeds the area covered by any one class of plant growth-form. They are named from the leading plant species when plant cover is $\geq 1\%$.
- Boulder field:** Land in which the area of unconsolidated boulders (> 200 mm diam.) exceeds the area covered by any one class of plant growth-form. Boulder fields are named from the leading plant species when plant cover is $\geq 1\%$.
- Cobble field:** Land in which the area of unconsolidated cobbles (20–200 mm diam.) exceeds the area covered by any one class of plant growth-form. Cobble fields are named from the leading plant species when plant cover is $\geq 1\%$.
- Gravel field:** Land in which the area of unconsolidated gravel (2–20 mm diameter) exceeds the area covered by any one class of plant growth-form. Gravel fields are named from the leading plant species when plant cover is $\geq 1\%$.
- Mobile sand:** Granular beach sand characterised by a rippled surface layer from strong tidal or wind-generated currents. Often forms bars and beaches.
- Firm or soft sand:** Sand flats may be mud-like in appearance but are granular when rubbed between the fingers and no conspicuous fines are evident when sediment is disturbed e.g. a mud content $< 1\%$. Classified as firm sand if an adult sinks < 2 cm or soft sand if an adult sinks > 2 cm.
- Firm muddy sand:** A sand/mud mixture dominated by sand with a moderate mud fraction (e.g. 1–10%), the mud fraction conspicuous only when sediment is mixed in water. The sediment appears brown, and may have a black anaerobic layer below. From a distance appears visually similar to firm sandy mud, firm or soft mud, and very soft mud. When walking you'll sink 0–2 cm. Granular when rubbed between the fingers.
- Firm sandy mud:** A sand/mud mixture dominated by sand with an elevated mud fraction (e.g. 10–25%), the mud fraction visually conspicuous when walking on it. The surface appears brown, and may have a black anaerobic layer below. From a distance appears visually similar to firm muddy sand, firm or soft mud, and very soft mud. When walking you'll sink 0–2 cm. Granular when rubbed between the fingers, but with a smoother consistency than firm muddy sand.
- Firm or soft mud:** A mixture of mud and sand where mud is a major component (e.g. $> 25\%$ mud). Sediment rubbed between the fingers retains a granular component but is primarily smooth/silken. The surface appears grey or brown, and may have a black anaerobic layer below. From a distance appears visually similar to firm muddy sand, firm sandy mud, and very soft mud. Classified as firm mud if an adult sinks < 5 cm (usually if sediments are dried out or another component e.g. gravel prevents sinking) or soft mud if an adult sinks > 5 cm.
- Very soft mud:** A mixture of mud and sand where mud is the major component (e.g. $> 50\%$ mud), the surface appears brown, and may have a black anaerobic layer below. When walking you'll sink > 5 cm unless another component e.g. gravel prevents sinking. From a distance appears visually similar to firm muddy sand, firm sandy mud, and firm or soft mud. Sediment rubbed between the fingers may retain a slight granular component but is primarily smooth/silken.
- Cockle bed /Mussel reef/ Oyster reef:** Area that is dominated by both live and dead cockle shells, or one or more mussel or oyster species respectively.
- Sabellid field:** Area that is dominated by raised beds of sabellid polychaete tubes.
- Shell bank:** Area that is dominated by dead shells.
- Artificial structures:** Introduced natural or man-made materials that modify the environment. Includes rip-rap, rock walls, wharf piles, bridge supports, walkways, boat ramps, sand replenishment, groynes, flood control banks, stopgates.

APPENDIX 2. NOTES SUPPORTING RISK INDICATOR RATINGS (TABLE 2)

NOTES to Table 2: See Robertson et al. (2016a, 2016b) for further information supporting these ratings.

Soft Mud Percent Cover. Soft mud (>25% mud content) has been shown to result in a degraded macroinvertebrate community (Robertson et al. 2015, 2016), and excessive mud decreases water clarity, lowers biodiversity and affects aesthetics and access. Because estuaries are a sink for sediments, the presence of large areas of soft mud is likely to lead to major and detrimental ecological changes that could be very difficult to reverse. In particular, its presence indicates where changes in land management may be needed. If an estuary is suspected of being an outlier (e.g. has >25% mud content but substrate remains firm to walk on), it is recommended that the initial broad scale assessment be followed by particle grain size analyses of relevant areas to determine the extent of the estuary with sediment mud contents >25%.

Sedimentation Mud Content. Below mud contents of 20–30% sediments are relatively incohesive and firm to walk on. Above this, they become sticky and cohesive and are associated with a significant shift in the macroinvertebrate assemblage to a lower diversity community tolerant of muds. This is particularly pronounced if elevated mud contents are contiguous with elevated total organic carbon concentrations, which typically increase with mud content, as do the concentrations of sediment bound nutrients and heavy metals. Consequently, muddy sediments are often poorly oxygenated, nutrient rich, and on intertidal flats of estuaries can be overlain with dense opportunistic macroalgal blooms. High mud contents also contribute to poor water clarity through ready resuspension of fine muds, impacting on seagrass, birds, fish and aesthetic values.

apparent Redox Potential Discontinuity (aRPD). aRPD depth, the transition between oxygenated sediments near the surface and deeper anoxic sediments, is a primary estuary condition indicator as it is a direct measure of whether nutrient and organic enrichment exceeds levels causing nuisance (anoxic) conditions. Knowing if the aRPD is close to the surface is important for two main reasons:

1. As the aRPD layer gets close to the surface, a “tipping point” is reached where the pool of sediment nutrients (which can be large), suddenly becomes available to fuel algal blooms and to worsen sediment conditions.
2. Anoxic sediments contain toxic sulphides and support very little aquatic life.

In sandy porous sediments, the aRPD layer is usually relatively deep (>3cm) and is maintained primarily by current or wave action that pumps oxygenated water into the sediments. In finer silt/clay sediments, physical diffusion limits oxygen penetration to <1cm (Jørgensen and Revsbech 1985) unless bioturbation by infauna oxygenates the sediments. The tendency for sediments to become anoxic is much greater if the sediments are muddy.

Redox Potential (Eh). For meter approaches, Eh measurements represent a composite of multiple redox equilibria measured at the surface of a redox potential electrode coupled to a millivolt meter (Rosenberg et al. 2001) (often called an ORP meter) and reflects a system’s tendency to receive or donate electrons. The electrode is inserted to different depths into the sediment and the extent of reducing conditions at each depth recorded (RPD is the depth at which the redox potential is ~0mV, Fenchel and Riedl 1970, Revsbech et al. 1980, Birchenough et al. 2012, Hunting et al. 2012). The Eh rating bands reflect the presence of healthy macrofauna communities in sediments below the aRPD depth.

Opportunistic Macroalgae. The presence of opportunistic macroalgae is a primary indicator of estuary eutrophication, and when combined with gross eutrophic conditions (see previous) can cause significant adverse ecological impacts that are very difficult to reverse. Thresholds used to assess this indicator are derived from the OMBT (see Section 3 and Appendix 2), with results combined with those of other indicators to determine overall condition.

Seagrass. Seagrass (*Zostera muelleri*) grows in soft sediments in most NZ estuaries. It is widely acknowledged that the presence of healthy seagrass beds enhances estuary biodiversity and particularly improves benthic ecology (Nelson 2009). Though tolerant of a wide range of conditions, it is seldom found above mean sea level (MSL), and is vulnerable to fine sediments in the water column and sediment quality (particularly if there is a lack of oxygen and production of sulphide), rapid sediment deposition, excessive macroalgal growth, high nutrient concentrations, and reclamation. Decreases in seagrass extent is likely to indicate an increase in these types of pressures.

As a baseline measure of seagrass presence, a continuous index (the seagrass coefficient - SC) has been developed to rate seagrass condition based on the percentage cover of seagrass in defined categories using the following equation: $SC = ((0 \times \% \text{seagrass cover} < 1\%) + (0.5 \times \% \text{cover } 1-5\%) + (2 \times \% \text{cover } 6-10\%) + (3.5 \times \% \text{cover } 11-20\%) + (6 \times \% \text{cover } 21-50\%) + (9 \times \% \text{cover } 51-80\%) + (12 \times \% \text{cover } > 80\%))/100$. Because estuaries are likely to support variable natural seagrass extents, the SC rating is intended to highlight estuaries with low seagrass cover for further evaluation (i.e. estimate natural seagrass cover to determine current state), and to provide an estuary specific metric against which future change can be assessed. It is not intended that the SC be used to directly compare different estuaries. The “early warning trigger” for initiating management action is a trend of decreasing SC.

Saltmarsh. Saltmarshes have high biodiversity, are amongst the most productive habitats on earth, and have strong aesthetic appeal. They are sensitive to a wide range of pressures including land reclamation, margin development, flow regulation, sea level rise, grazing, wastewater contaminants, and weed invasion. Most NZ estuarine saltmarsh grows in the upper estuary margins above mean high water neap (MHWN) tide where vegetation stabilises fine sediment transported by tidal flows. Saltmarsh zonation is commonly evident, resulting from the combined influence of factors including salinity, inundation period, elevation, wave exposure, and sediment type. Highest saltmarsh diversity is generally present above mean high water spring (MHWS) tide where a variety of salt tolerant species grow including scrub, sedge, tussock, grass, reed, rush and herb fields. Between MHWS and MHWN, saltmarsh is commonly dominated by relatively low diversity rushland and herbfields. Below this, the MHWN to MSL range is commonly unvegetated or limited to either mangroves or *Spartina*, the latter being able to grow to MLWN. Further work is required to develop a comprehensive saltmarsh metric for NZ. As an interim measure, the % of the intertidal area comprising saltmarsh is used to indicate saltmarsh condition. Two supporting metrics are also proposed: i. % loss from Estimated Natural State Cover. This assumes that a reduction in natural state saltmarsh cover corresponds to a reduction in ecological services and habitat values. ii. % of available habitat supporting saltmarsh. This assumes that saltmarsh should be growing throughout the majority of the available saltmarsh habitat (tidal area above MHWN), and that where this does not occur, ecological services and habitat values are reduced. The interim risk ratings proposed for these ratings are Very Low=>80-100%, Low=>60-80%, Moderate=>40-60%, and High=<40%. The “early warning trigger” for initiating management action/further investigation is a trend of a decreasing saltmarsh area or saltmarsh growing over <80% of the available habitat.

Vegetated Margin. The presence of a terrestrial margin dominated by a dense assemblage of scrub/shrub and forest vegetation acts as an important buffer between developed areas and the saltmarsh and estuary. This buffer is sensitive to a wide range of pressures including land reclamation, margin development, flow regulation, sea level rise, grazing, wastewater contaminants, and weed invasion. It protects the estuary against introduced weeds and grasses, naturally filters sediments and nutrients, and provides valuable ecological habitat. Reduction in the vegetated terrestrial buffer around the estuary is likely to result in a decline in estuary quality. The “early warning trigger” for initiating management action is <50% of the estuary with a densely vegetated margin.

Change from Baseline Condition. Where natural state conditions for high value habitat of seagrass, saltmarsh, and densely vegetated terrestrial margin are unknown it is proposed that % change from the first measured baseline condition be used to determine trends in estuary condition. It is assumed that increases in such habitat are desirable (i.e. represent a Very Low risk rating), and decreases are undesirable. For decreases, the interim risk ratings proposed are: Very Low=<5%, Low=>5-10%, Moderate=>10-20%, and High=>20%. For indicators of degraded habitat e.g. extent of soft mud or gross eutrophic conditions, the same interim risk rating bands are proposed, but are applied to increases in extent.

APPENDIX 3. NOTES ON SAMPLING, RESOLUTION AND ACCURACY

Sediment sampling and analysis

Grain size samples were collected from representative mud and sand habitats (to validate substrate classifications) by sampling a composite of the top 20mm of sediment (approx. 250gms in total) using a plastic trowel. Samples were placed inside a numbered plastic bag, refrigerated within 4 hours of sample collection before being frozen and sent to R.J. Hill Laboratories for grain size analysis (% mud, sand, gravel). Details of lab methods and detection limits are presented in Appendix 6. Samples were tracked using standard Chain of Custody forms and results were checked and transferred electronically to avoid transcription errors.

In addition, at selected sampling sites redox potential (RP) was measured with an oxidation-reduction potential meter at 0, 1, 3, 6 and 10cm depths below the substrate surface, and the aRPD depth and substrate type recorded. These results have been used to generate broad scale maps showing areas where sediment oxygenation is depleted to the extent that adverse impacts to macrofauna (sediment and surface dwelling animals) are expected i.e. where RPD at 3cm <-150mV or aRPD <1cm (Robertson et al. 2016b).

Sampling resolution and accuracy

Estimates of error for different measurements have been made based on the field data collected to date. Initial broad scale mapping is intended to provide a rapid overview of estuary condition based on the mapping of features visible on aerial photographs, supported by ground-truthing to validate the visible features. The accuracy of mapping is therefore primarily determined by the resolution of the available photos, and secondarily by the extent of groundtruthing. In most instances features with readily defined edges such as saltmarsh beds, rockfields etc. can be accurately mapped to within 1-2m of their boundaries. The largest area for potential error is where boundaries are not readily visible on photographs e.g. where firm muddy sands transition to soft muds. These boundaries require field validation. Extensive mapping experience has shown that it is possible to define such boundaries to within $\pm 10\text{m}$ where they have been thoroughly ground-truthed using NEMP classifications. Because broad scale mapping necessitates the grouping of variable and non-uniform patches (which introduces a certain amount of variation) overall broad scale accuracy is unlikely to exceed $\pm 10\%$ for boundaries not readily visible on photographs.

Where initial broad scale mapping results indicate a need for greater resolution of boundaries (e.g. to increase certainty about the extent of soft mud areas), or to define changes within NEMP categories (e.g. to define the mud content within firm muddy sand habitat), then issue-specific approaches are recommended. The former includes more widespread ground-truthing, and the latter uses transect or grid based grain size sampling.

For specific broad scale seagrass and macroalgae features that are spatially and temporally variable, the overall spatial extent, and boundaries between different percentage cover and density areas, are considered accurate to within $\pm 10\text{m}$ where they have been thoroughly ground-truthed using NEMP classifications. Accuracy declines when assessed remotely e.g. from aerial photographs, and particularly so when assessing lower density (<50%) cover which is commonly not visible on aerial coverages. As previously, the most accurate measures are obtained with increasing field time (and cost).

Within mapped boundaries, broad scale estimates of percentage cover and density, due to the grouping of variable and non-uniform patches, are considered accurate to $\pm 10\%$. These however can be assessed to a much higher degree of accuracy using fine scale quadrat based approaches such as the OMBT which can also be increased by applying fine scale approaches estuary-wide if a very high degree of accuracy is considered important.

For the OMBT, a methodology for calculating a measure of the confidence of class (CofC), has been developed (Davey, 2009) that defines the specific accuracy of the measures undertaken. Called CAP-TAIN ('Confidence And Precision Tool Aids aNalysis') it calculates CofC at three levels: i. metric, ii. survey (single sampling event), and iii. water body over the reporting period (potentially several surveys). The upper and lower 90% Confidence Intervals for the SE of the EQR are presented in this report.

APPENDIX 4. OPPORTUNISTIC MACROALGAL BLOOMING TOOL

The UK-WFD (Water Framework Directive) Opportunistic Macroalgal Blooming Tool (OMBT) (WFD-UKTAG 2014) is a comprehensive 5 part multimetric index approach suitable for characterising the different types of estuaries and related macroalgal issues found in NZ. The tool allows simple adjustment of underpinning threshold values to calibrate it to the observed relationships between macroalgal condition and the ecological response of different estuary types. It incorporates sediment entrained macroalgae, a key indicator of estuary degradation, and addresses limitations associated with percentage cover estimates that do not incorporate biomass e.g. where high cover but low biomass are not resulting in significantly degraded sediment conditions. It is supported by extensive studies of the macroalgal condition in relation to ecological responses in a wide range of estuaries.

The 5 part multimetric OMBT, modified for NZ estuary types, is fully described below. It is based on macroalgal growth within the Available Intertidal Habitat (AIH) - the estuary area between high and low water spring tide able to support opportunistic macroalgal growth. Suitable areas are considered to consist of *mud, muddy sand, sandy mud, sand, stony mud and mussel beds*. Areas which are judged unsuitable for algal blooms e.g. channels and channel edges subject to constant scouring, need to be excluded from the AIH. The following measures are then taken:

1. Percentage cover of the available intertidal habitat (AIH).

The percent cover of opportunistic macroalgal within the AIH is assessed. While a range of methods are described, visual rating by experienced ecologists, with independent validation of results is a reliable and rapid method. All areas within the AIH with macroalgal cover >5% are mapped spatially.

2. Total extent of area covered by algal mats (affected area (AA)) or affected area as a percentage of the AIH (AA/AIH, %).

In large water bodies with proportionately small patches of macroalgal coverage, the rating for total area covered by macroalgae (Affected Area - AA) might indicate high or good status, while the total area covered could actually be quite substantial and could still affect the surrounding and underlying communities. In order to account for this, an additional metric established is the affected area as a percentage of the AIH (i.e. $(AA/AIH) \times 100$). This helps to scale the area of impact to the size of the water body. In the final assessment the lower of the two metrics (the AA or percentage AA/AIH) is used, i.e. whichever reflects the worse case scenario.

3. Biomass of AIH ($g \cdot m^{-2}$).

Assessment of the spatial extent of the algal bed alone will not indicate the level of risk to a water body. For example, a very thin (low biomass) layer covering over 75% of a shore might have little impact on underlying sediments and fauna. The influence of biomass is therefore incorporated. Biomass is calculated as a mean for (i) the whole of the AIH and (ii) for the Affected Areas. The potential use of maximum biomass was rejected, as it could falsely classify a water body by giving undue weighting to a small, localised blooming problem. Algae growing on the surface of the sediment are collected for biomass assessment, thoroughly rinsed to remove sediment and invertebrate fauna, hand squeezed until water stops running, and the wet weight of algae recorded.

For quality assurance of the percentage cover estimates, two independent readings should be within +/- 5%. A photograph should be taken of every quadrat for inter-calibration and cross-checking of percent cover determination. Measures of biomass should be calculated to 1 decimal place of wet weight of sample. For both procedures the accuracy should be demonstrated with the use of quality assurance checks and procedures.

4. Biomass of AA ($g \cdot m^{-2}$).

Mean biomass of the Affected Area (AA), with the AA defined as the total area with macroalgal cover >5%.

5. Presence of Entrained Algae (percentage of quadrats).

Algae are considered as entrained in muddy sediment when they are found growing >3cm deep within muddy sediments. The persistence of algae within sediments provides both a means for over-wintering of algal spores and a source of nutrients within the sediments. Build-up of weed within sediments therefore implies that blooms can become self-regenerating given the right conditions (Raffaelli et al. 1989). Absence of weed within the sediments lessens the likelihood of bloom persistence, while its presence gives greater opportunity for nutrient exchange with sediments. Consequently, the presence of opportunistic macroalgae growing within the surface sediment was included in the tool.

All the metrics are equally weighted and combined within the multimetric, in order to best describe the changes in the nature and degree of opportunist macroalgal growth on sedimentary shores due to nutrient pressure.

Timing: Because the OMBT has been developed to classify data over the maximum growing season, sampling should target the peak bloom in summer (Dec-March), although peak timing may vary among water bodies, therefore local knowledge is required to identify the maximum growth period. Sampling is not recommended outside the summer period due to seasonal variations that could affect the outcome of the tool and possibly lead to misclassification; e.g. blooms may become disrupted by stormy autumn weather and often die back in winter. Sampling should be carried out during spring low tides in order to access the maximum area of the AIH.

APPENDIX 4. OPPORTUNISTIC MACROALGAL BLOOMING TOOL

Suitable Locations: The OMBT is suitable for use in estuaries and coastal waters which have intertidal areas of soft sedimentary substratum (i.e. areas of AIH for opportunistic macroalgal growth). The tool is not currently used for assessing ICOLLs due to the particular challenges in setting suitable reference conditions for these water bodies.

Derivation of Threshold Values.

Published and unpublished literature, along with expert opinion, was used to derive critical threshold values suitable for defining quality status classes (Table A2).

- **Reference Thresholds.** A UK Department of the Environment, Transport and the Regions (DETR) expert workshop suggested reference levels of <5% cover of AIH of climax and opportunistic species for high quality sites (DETR, 2001). In line with this approach, the WFD adopted <5% cover of opportunistic macroalgae in the AIH as equivalent to High status. From the WFD North East Atlantic inter-calibration phase 1 results, German research into large sized water bodies revealed that areas over 50ha may often show signs of adverse effects, however if the overall area was less than 1/5th of this adverse effects were not seen, so the High/Good boundary was set at 10ha. In all cases a reference of 0% cover for truly un-impacted areas was assumed. Note: opportunistic algae may occur even in pristine water bodies as part of the natural community functioning.

The proposal of reference conditions for levels of biomass took a similar approach, considering existing guidelines and suggestions from DETR (2001), with a tentative reference level of <100g m⁻² wet weight. This reference level was used for both the average biomass over the affected area and the average biomass over the AIH. As with area measurements a reference of zero was assumed.

An ideal of no entrainment (i.e. no quadrats revealing entrained macroalgae) was assumed to be reference for un-impacted waters. After some empirical testing in a number of UK water bodies a High / Good boundary of 1% of quadrats was set.

- **Class Thresholds for Percent Cover:**

High/Good boundary set at 5%. Based on the finding that a symptom of the potential start of eutrophication is when: (i) 25% of the available intertidal habitat has opportunistic macroalgae and (ii) at least 25% of the sediment (i.e. 25% in a quadrat) is covered (Comprehensive Studies Task Team (DETR, 2001)). This implies that an overall cover of the AIH of 6.25% (25*25%) represents the start of a potential problem.

Good / Moderate boundary set at 15%. True problem areas often have a >60% cover within the affected area of 25% of the water body (Wither 2003). This equates to 15% overall cover of the AIH (i.e. 25% of the water body covered with algal mats at a density of 60%). **Poor/Bad boundary** is set at >75%. The Environment Agency has considered >75% cover as seriously affecting an area (Foden et al. 2010).

- **Class Thresholds for Biomass.** Class boundaries for biomass values were derived from DETR (2001) recommendations that <500 g.m⁻² wet weight was an acceptable level above the reference level of <100 g.m⁻² wet weight. In Good status only slight deviation from High status is permitted so 500 g.m⁻² represents the Good/Moderate boundary. Moderate quality status requires moderate signs of distortion and significantly greater deviation from High status to be observed. The presence of >500 g.m⁻² but less than 1,000 g.m⁻² would lead to a classification of Moderate quality status at best, but would depend on the percentage of the AIH covered. >1kg.m⁻² wet weight causes significant harmful effects on biota (DETR 2001, Lowthion et al. 1985, Hull 1987, Wither 2003).
- **Thresholds for Entrained Algae.** Empirical studies testing a number of scales were undertaken on a number of impacted waters. Seriously impacted waters have a very high percentage (>75%) of the beds showing entrainment (Poor / Bad boundary). Entrainment was felt to be an early warning sign of potential eutrophication problems so a tight High /Good standard of 1% was selected (this allows for the odd change in a quadrat or error to be taken into account). Consequently the Good / Moderate boundary was set at 5% where (assuming sufficient quadrats were taken) it would be clear that entrainment and potential over wintering of macroalgae had started.

Each metric in the OMBT has equal weighting and is combined to produce the ecological quality ratio score (EQR).

Table A2. The final face value thresholds and metrics for levels of ecological quality status in the UK-WFD 2014.

Quality Status	High	Good	Moderate	Poor	Bad
EQR (Ecological Quality Rating)	≥0.8 - 1.0	≥0.6 - <0.8	≥0.4 - <0.6	≥0.2 - <0.4	0.0 - <0.2
% cover on Available Intertidal Habitat (AIH)	0 - ≤5	>5 - ≤15	>15 - ≤25	>25 - ≤75	>75 - 100
Affected Area (AA) of >5% macroalgae (ha)*	≥0 - 10	≥10 - 50	≥50 - 100	≥100 - 250	≥250
AA/AIH (%)*	≥0 - 5	≥5 - 15	≥15 - 50	≥50 - 75	≥75 - 100
Average biomass (g.m ²) of AIH	≥0 - 100	≥100 - 500	≥500 - 1000	≥1000 - 3000	≥3000
Average biomass (g.m ²) of AA	≥0 - 100	≥100 - 500	≥500 - 1000	≥1000 - 3000	≥3000
% algae >3cm deep	≥0 - 1	≥1 - 5	≥5 - 20	≥20 - 50	≥50 - 100

*N.B. Only the lower EQR of the 2 metrics, AA or AA/AIH is used in the final EQR calculation.

APPENDIX 4. OPPORTUNISTIC MACROALGAL BLOOMING TOOL

EQR calculation

Each metric in the OMBT has equal weighting and is combined to produce the **Ecological Quality Ratio** score (EQR).

The face value metrics work on a sliding scale to enable an accurate metric EQR value to be calculated; an average of these values is then used to establish the final water body level EQR and classification status. The EQR determining the final water body classification ranges between a value of zero to one and is converted to a Quality Status by using the following categories:

Quality Status	High	Good	Moderate	Poor	Bad
EQR (Ecological Quality Rating)	≥0.8 - 1.0	≥0.6 - <0.8	≥0.4 - <0.6	≥0.2 - <0.4	0.0 - <0.2

The EQR calculation process is as follows:

1. Calculation of the face value (e.g. percentage cover of AIH) for each metric. To calculate the individual metric face values:

- Percentage cover of AIH (%) = (Total % Cover / AIH) x 100 - where Total % cover = Sum of {(patch size) / 100} x average % cover for patch
- Affected Area, AA (ha) = Sum of all patch sizes (with macroalgal cover >5%).
- Biomass of AIH (g.m⁻²) = Total biomass / AIH - where Total biomass = Sum of (patch size x average biomass for the patch)
- Biomass of Affected Area (g.m⁻²) = Total biomass / AA - where Total biomass = Sum of (patch size x average biomass for the patch)
- Presence of Entrained Algae = (No. quadrats with entrained algae / total no. of quadrats) x 100
- Size of AA in relation to AIH (%) = (AA/AIH) x 100

2. Normalisation and rescaling to convert the face value to an equidistant index score (0-1 value) for each index (Table A3).

The face values are converted to an equidistant EQR scale to allow combination of the metrics. These steps have been mathematically combined in the following equation:

$$\text{Final Equidistant Index score} = \text{Upper Equidistant range value} - \left(\frac{\text{Face Value} - \text{Upper Face value range}}{\text{Equidistant class range} / \text{Face Value Class Range}} \right) *$$

Table A3 gives the critical values at each class range required for the above equation. The first three numeric columns contain the face values (FV) for the range of the index in question, the last three numeric columns contain the values of the equidistant 0-1 scale and are the same for each index. The face value class range is derived by subtracting the upper face value of the range from the lower face value of the range.

Note: the table is “simplified” with rounded numbers for display purposes. The face values in each class band may have greater than (>) or less than (<) symbols associated with them, for calculation a value of <5 is given a value of 4.999’.

The final EQR score is calculated as the average of equidistant metric scores.

A spreadsheet calculator is available to download from the UK WFD website to undertake the calculation of EQR scores.

References

- DETR, 2001. *Development of ecological quality objectives with regard to eutrophication. Final report, unpublished.*
- Foden, J., Wells, E., Scanlan, C. and Best M.A. 2010. *Water Framework Directive development of classification tools for ecological assessment: Opportunistic Macroalgae Blooming. UK TAG Report for Marine Plants Task Team, January 2010, Publ. UK TAG.*
- Hull, S.C. 1987. *Macroalgal mats and species abundance: a field experiment. Estuar. Coast. Shelf Sci. 25, 519-532.*
- Lowthion, D., Soulsby, P.G. and Houston, M.C.M. 1985. *Investigation of a eutrophic tidal basin: 1. Factors affecting the distribution and biomass of macroalgae. Marine Environmental Research 15: 263-284.*
- Raffaelli, D., Hull, S. and Milne, H. 1989. *Long-term changes in nutrients, weedmats and shore birds in an estuarine system. Cah. Biol. Mar. 30, 259-270.*
- WFD-UKTAG (Water Framework Directive – United Kingdom Technical Advisory Group) 2014. *UKTAG Transitional and Coastal Water Assessment Method Macroalgae Opportunistic Macroalgal Blooming Tool. Retrieved from [http://www.wfduk.org/sites/default/files/Media/Characterisation of the water environment/Biological Method Statements/TraC Macroalgae OMBT UKTAG Method Statement.PDF](http://www.wfduk.org/sites/default/files/Media/Characterisation%20of%20the%20water%20environment/Biological%20Method%20Statements/TraC%20Macroalgae%20OMBT%20UKTAG%20Method%20Statement.PDF).*
- Wither, A. 2003. *Guidance for sites potentially impacted by algal mats (green seaweed). EC Habitats Directive Technical Advisory Group report WQTAG07c.*

APPENDIX 4. OPPORTUNISTIC MACROALGAL BLOOMING TOOL

Table A3. Values for the normalisation and re-scaling of face values to EQR metric.

METRIC	QUALITY STATUS	FACE VALUE RANGES			EQUIDISTANT CLASS RANGE VALUES		
		Lower face value range (measurements towards the "Bad" end of this class range)	Upper face value range (measurements towards the "High" end of this class range)	Face Value Class Range	Lower 0-1 Equidistant range value	Upper 0-1 Equidistant range value	Equidistant Class Range
% Cover of Available Intertidal Habitat (AIH)	High	≤5	0	5	≥0.8	1	0.2
	Good	≤15	>5	9.999	≥0.6	<0.8	0.2
	Moderate	≤25	>15	9.999	≥0.4	<0.6	0.2
	Poor	≤75	>25	49.999	≥0.2	<0.4	0.2
	Bad	100	>75	24.999	0	<0.2	0.2
Average Biomass of AIH (g m ⁻²)	High	≤100	0	100	≥0.8	1	0.2
	Good	≤500	>100	399.999	≥0.6	<0.8	0.2
	Moderate	≤1000	>500	499.999	≥0.4	<0.6	0.2
	Poor	≤3000	>1000	1999.999	≥0.2	<0.4	0.2
	Bad	≤6000	>3000	2999.999	0	<0.2	0.2
Average Biomass of Affected Area (AA) (g m ⁻²)	High	≤100	0	100	≥0.8	1	0.2
	Good	≤500	>100	399.999	≥0.6	<0.8	0.2
	Moderate	≤1000	>500	499.999	≥0.4	<0.6	0.2
	Poor	≤3000	>1000	1999.999	≥0.2	<0.4	0.2
	Bad	≤6000	>3000	2999.999	0	<0.2	0.2
Affected Area (Ha)*	High	≤10	0	100	≥0.8	1	0.2
	Good	≤50	>10	39.999	≥0.6	<0.8	0.2
	Moderate	≤100	>50	49.999	≥0.4	<0.6	0.2
	Poor	≤250	>100	149.999	≥0.2	<0.4	0.2
	Bad	≤6000	>250	5749.999	0	<0.2	0.2
AA/AIH (%)*	High	≤5	0	5	≥0.8	1	0.2
	Good	≤15	>5	9.999	≥0.6	<0.8	0.2
	Moderate	≤50	>15	34.999	≥0.4	<0.6	0.2
	Poor	≤75	>50	24.999	≥0.2	<0.4	0.2
	Bad	100	>75	27.999	0	<0.2	0.2
% Entrained Algae	High	≤1	0	1	≥0.0	1	0.2
	Good	≤5	>1	3.999	≥0.2	<0.0	0.2
	Moderate	≤20	>5	14.999	≥0.4	<0.2	0.2
	Poor	≤50	>20	29.999	≥0.6	<0.4	0.2
	Bad	100	>50	49.999	1	<0.6	0.2

*N.B. Only the lower EQR of the 2 metrics, AA or AA/AIH should be used in the final EQR calculation.

Table A4. The final face value thresholds and metrics for levels of ecological quality status used to rate opportunistic macroalgae in the current in the study (modified from UK-WFD 2014).

MACROALGAL INDICATORS (OBMT approach - WFD_UKTAG 2014)					
QUALITY RATING	High	Good	Moderate	Poor	Bad
EQR (Ecological Quality Rating)	≥0.8 - 1.0	≥0.6 - <0.8	≥0.4 - <0.6	≥0.2 - <0.4	0.0 - <0.2
% cover on Available Intertidal Habitat (AIH)	0 - ≤5	>5 - ≤15	>15 - ≤25	>25 - ≤75	>75 - 100
Affected Area (AA) [>5% macroalgae] (ha)*	≥0 - 10	≥10 - 50	≥50 - 100	≥100 - 250	≥250
AA/AIH (%)*	≥0 - 5	≥5 - 15	≥15 - 50	≥50 - 75	≥75 - 100
Average biomass (g.m ² wet wgt) of AIH	≥0 - 100	≥100 - 200	≥200 - 500	≥500 - 1450	≥1450
Average biomass (g.m ² wet wgt) of AA	≥0 - 100	≥100 - 200	≥200 - 500	≥500 - 1450	≥1450
% algae entrained >3cm deep	≥0 - 1	≥1 - 5	≥5 - 20	≥20 - 50	≥50 - 100

*Only the lower EQR of the 2 metrics, AA or AA/AIH is used in the final EQR calculation.

APPENDIX 5. WAIKOUAITI ESTUARY MACROALGAL DATA




Figure A1. Location of macroalgal patches (>5% cover) used in assessing Waikouaiti Estuary, Dec 2016.

APPENDIX 5. WAIKOUAITI ESTUARY MACROALGAL DATA

Macroalgal cover >15% used in calculating the OMBT EQR (see Figure A1 for locations).

Patch ID	Patch area (ha)	Quadrat No	Percent cover of macroalgae	Mean Biomass (g.m ⁻² wet weight)	Presence (1) or absence (0) of entrained algae	aRPD depth (cm)	Presence (1) or absence (0) of soft mud	Dominant species
1	0.47	0	80	300	0	3	0	<i>Ulva lactuca</i> <i>Gracilaria chilensis</i>
2	1.17	0	50	300	0	3	0	<i>Ulva lactuca</i> <i>Gracilaria chilensis</i>
3	0.84	0	15	50	0	3	0	<i>Ulva lactuca</i>
0	0.78	0	10	50	0	1	0	<i>Ulva lactuca</i> <i>Gracilaria chilensis</i>
4	5.98	0	15	50	0	1	0	<i>Ulva lactuca</i>
5	2.96	0	90	2500	0	1	0	<i>Ulva lactuca</i>
6	0.54	0	80	300	0	>5	0	<i>Ulva lactuca</i>
7	0.03	0	80	400	0	1	1	<i>Ulva lactuca</i>
8	4.52	0	35	500	0	1	0	<i>Ulva lactuca</i> <i>Gracilaria chilensis</i>
9	1.28	0	20	300	0	1	0	<i>Ulva lactuca</i> <i>Gracilaria chilensis</i>
10	1.57	0	15	100	0	>5	0	<i>Ulva lactuca</i>
11	1.23	0	10	50	0	1	0	<i>Ulva lactuca</i>
12	0.47	0	35	300	0	>5	0	<i>Ulva lactuca</i>

APPENDIX 6. DETAILS OF ANALYTICAL METHODS

Sediment Indicator	Laboratory	Method	Detection Limit
Grain Size	R.J Hill	Wet sieving, gravimetric (calculation by difference)	0.1 g/100g dry wgt